




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 227/08, 247/14, C07D 317/46, 203/26, 303/40, C07F 7/18, C07C 227/16		A1	(11) International Publication Number: WO 98/07685
			(43) International Publication Date: 26 February 1998 (26.02.98)
(21) International Application Number: PCT/US97/14813		<p>MUNGER, John, D. [US/US]; 1044 Catherine Street, Alviso, CA 95002 (US). PRISBE, Ernest, J. [US/US]; 1336 Richardson Avenue, Los Altos, CA 94024 (US). POSTICH, Michael, J. [US/US]; 2715 S. Norfolk Street #109, San Mateo, CA 94403 (US). ROHLOFF, John, C. [US/US]; 1654 Cornell Drive, Mountain View, CA 94040 (US). ST. JOHN, Daphne, E. [US/US]; 1404 33rd Avenue, San Francisco, CA 94122 (US). WILLIAMS, Matthew, A. [US/US]; 1077 Foster City Boulevard #C, Foster City, CA 94404 (US). ZHANG, Lijun [CN/US]; 845 Shell Boulevard #203L, Foster City, CA 94404 (US).</p> <p>(74) Agents: BOSSE, Mark, L. et al.; Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, CA 94404 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	
(22) International Filing Date: 22 August 1997 (22.08.97)			
(30) Priority Data: 60/024,122 23 August 1996 (23.08.96) US 08/701,942 23 August 1996 (23.08.96) US			
(60) Parent Applications or Grants (63) Related by Continuation US 60/024,122 (CON) Filed on 23 August 1996 (23.08.96) US 08/701,942 (CON) Filed on 23 August 1996 (23.08.96)			
(71) Applicant (for all designated States except US): GILEAD SCIENCES, INC. [US/US]; 333 Lakeside Drive, Foster City, CA 94404 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): KENT, Kenneth, M. [US/US]; 565 Fir Avenue, Sunnyvale, CA 94086 (US). KIM, Choung, U. [US/US]; 1750 Elizabeth Street, San Carlos, CA 94070 (US). MCGEE, Lawrence, R. [US/US]; 1399 Terra Nova Boulevard, Pacifica, CA 94044 (US).			

(54) Title: PREPARATION OF CYCLOHEXENE CARBOXYLATE DERIVATIVES



09/555,442

(57) Abstract

The present invention provides new synthetic methods and compositions. In particular, new methods of preparing intermediates such as those having formulas (I)-(IV), such as those having formulas (I)-(IV), useful in the synthesis of neuraminidase inhibitors and compositions useful as intermediates that are themselves useful in the synthesis of neuraminidase inhibitors are provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5

PREPARATION OF CYCLOHEXENE CARBOXYLATE DERIVATIVES

10

Background of the InventionField of the Invention

15 The present invention is directed to methods of preparing carbocyclic compounds and intermediates therefore.

Brief Description of Related Art

United States Patent Application (having Attorney Docket No. 205.6) Serial Number 08/702,308, filed August 23, 1996, which was a
20 continuation-in-part application of United States Patent Application Serial Number 08/653,034, filed March 24, 1996, which was a continuation-in-part application of United States Patent Application Serial Number 08/606,624, filed February 26, 1996, which was a continuation-in-part application of United States Patent Application Serial Number 08/580,567, filed
25 December 29, 1995, which was a continuation-in-part application of United States Patent Application Serial Number 08/476,946, filed June 6, 1995, which was a continuation-in-part application of United States Patent Application Serial Number 08/395,245, filed February 27, 1995, all of which are incorporated herein by reference in their entirety, describe, inter alia,
30 neuraminadase inhibitors and intermediates in the synthesis of neuraminidase inhibitor. The present invention provides processes useful in the preparation of these compositions.

Objects of the Invention

35 Selected embodiments of the invention are directed to one or more of the following objects:

A principal object of the invention is to provide new synthetic

methods and compositions.

An additional object of the invention is to provide new methods of preparing intermediates useful in the synthesis of neuraminidase inhibitors.


- 5 An additional object of the invention is to provide compositions useful as intermediates that are themselves useful in the synthesis of neuraminidase inhibitors.

An additional object of the invention is to provide compositions useful as neuraminidase inhibitors.

10

Summary of the Invention

One aspect of the present invention is directed to processes for the preparation of compounds of the formula:



5

15 wherein:


R¹ is a cyclic hydroxy protecting group;

R² is a carboxylic acid protecting group;

R³ is a hydroxy protecting group; and

- 20 each R²⁰ is independently H or an alkyl of 1 to 12 carbon atoms;


which process comprises reaction of a compound of the formula:



4

with a dehydrating reagent.

Another aspect of the present invention is directed to processes for the preparation of compounds of the formula:



11

wherein:

- 5 each of R^2 , R^3 and R^{20} are as defined above;
 R^4 is $-C(R^{30})_3$;
each R^5 is independently H or R^3 ;
each R^7 is independently H or an amino protecting group;
each R^8 is independently H or R^2 ;
10 each R^9 is independently H or a thiol protecting group;
each R^{21} is independently R^{20} , Br, Cl, F, I, CN, NO_2 or N_3 ;
each R^{22} is independently F, Cl, Br, I, $-CN$, N_3 , $-NO_2$, $-OR^5$, $-OR^{20}$,
 $-N(R^{20})_2$, $-N(R^{20})(R^7)$, $-N(R^7)_2$, $-SR^{20}$, $-SR^9$, $-S(O)R^{20}$, $-S(O)_2R^{20}$, $-S(O)OR^{20}$,
 $-S(O)OR^8$, $-S(O)_2OR^{20}$, $-S(O)_2OR^8$, $-C(O)OR^{20}$, $-C(O)OR^8$, $-OC(O)R^{20}$,
15 $-N(R^{20})(C(O)R^{20})$, $-N(R^7)(C(O)R^{20})$, $-N(R^{20})(C(O)OR^{20})$, $-N(R^7)(C(O)OR^{20})$,
 $-C(O)N(R^{20})_2$, $-C(O)N(R^7)(R^{20})$, $-C(O)N(R^7)_2$, $-C(NR^{20})(N(R^{20})_2)$,
 $-C(N(R^7))(N(R^{20})_2)$, $-C(N(R^{20}))(N(R^{20})(R^7))$, $-C(N(R^7))(N(R^{20})(R^7))$,
 $-C(N(R^{20}))(N(R^7)_2)$, $-C(N(R^7))(N(R^7)_2)$, $-N(R^{20})C(N(R^{20}))(N(R^{20})_2)$,
 $-N(R^{20})C(N(R^{20}))(N(R^{20})(R^7))$, $-N(R^{20})C(N(R^7))(N(R^{20})_2)$,
20 $-N(R^7)C(N(R^{20}))(N(R^{20})_2)$, $-N(R^7)C(N(R^7))(N(R^{20})_2)$,
 $-N(R^7)C(N(R^{20}))(N(R^{20})(R^7))$, $-N(R^{20})C(N(R^7))(N(R^{20})(R^7))$,
 $-N(R^{20})C(N(R^{20}))(N(R^7)_2)$, $-N(R^7)C(N(R^7))(N(R^{20})(R^7))$,
 $-N(R^7)C(N(R^{20}))(N(R^7)_2)$, $-N(R^{20})C(N(R^7))(N(R^7)_2)$,
 $-N(R^7)C(N(R^7))(N(R^7)_2)$, $=O$, $=S$, $=N(R^{20})$, $=N(R^7)$ or W;
25 each R^{23} is independently alkyl of 1 to 11 carbon atoms, alkenyl of 2
to 11 carbon atoms, or alkynyl of 2 to 11 carbon atoms;
each R^{24} is independently R^{23} wherein each R^{23} is substituted with 0
to 3 R^{22} groups;
each R^{24a} is independently alkylene of 1 to 11 carbon atoms,
30 alkenylene of 2 to 11 carbon atoms, or alkynylene of 2-11 carbon atoms any

one of which alkylene, alkenylene or alkynylene is substituted with 0-3 R^{22} groups;


each R^{28} is independently alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12 carbon atoms, or alkynyl of 2 to 12 carbon atoms;

5 each R^{29} is independently R^{22} or R^{28} wherein each R^{28} is substituted with 0 to 3 R^{22} groups;

each R^{30} is independently H, R^{24} , W or $-R^{24a}W$; and

each W is independently carbocycle or heterocycle wherein any one of which carbocycle or heterocycle is substituted with 0 to 3 R^{29} groups;

10 which process comprises reaction of a compound of the formula:



10

wherein R^{31} is a ketal or acetal, with a lewis acid reagent;


provided that R^4 , taken as a whole, contains:

0 to 3 W groups substituted with 0 to 3 R^{29} groups; and, in

15 addition,

1 to 12 carbon atoms substituted with 0 to 3 R^{22} groups.

Another aspect of the present invention is directed to processes for the preparation of compounds of the formula:




31

20 wherein:

R^2 , R^4 , R^7 , R^{20} and R^{21} are as defined above.


which process comprises reaction of a compound of the formula:



30

with a reducing reagent.

Another aspect of the present invention is directed to processes for
5 the preparation of compounds of the formula:




41

wherein:

R^2 , R^4 , R^5 , R^{20} and R^{21} are as described above; and

Y^1 is a mono-, di- or unsubstituted amino group;


10 which process comprises reaction of a compound of the formula:



40

with an amine reagent.

Another aspect of the present invention is directed to processes for
15 the preparation of compounds of the formula:




51

wherein:

R^2 , R^4 , R^{20} , R^{21} and Y^1 are as described above;

which process comprises reaction of a compound of the formula:




50

5

with an oxidizing reagent.

Another aspect of the present invention is directed to processes for the preparation of compounds of the formula:




61

10

wherein:

R^2 , R^4 , R^{20} , R^{21} and Y^1 are as described above;


which process comprises reaction of a compound of the formula:



51

with a base.


Another aspect of the present invention is directed to processes for
5 the preparation of compounds of the formula:



71

wherein:

R², R⁴, R⁷, R²⁰, R²¹ and Y¹ are as described above;
which process comprises reaction of a compound of the formula:



61

with a reductive amination reagent.

Detailed Description

General

The present invention is directed to methods of making the
compositions described herein. Even though the compositions of the

invention are prepared by any of the applicable techniques of organic synthesis, the present invention provides advantageous methods for accomplishing the preparations.

Many conventional techniques are well known in the art and will not be elaborated here. However, many of the known techniques are elaborated in "Compendium of Organic Synthetic Methods" (John Wiley & Sons, New York), Vol. 1, Ian T. Harrison and Shuyen Harrison, 1971; Vol. 2, Ian T. Harrison and Shuyen Harrison, 1974; Vol. 3, Louis S. Hegedus and Leroy Wade, 1977; Vol. 4, Leroy G. Wade, jr., 1980; Vol. 5, Leroy G. Wade, Jr., 1984; and Vol. 6, Michael B. Smith; as well as March, J., "Advanced Organic Chemistry, Third Edition", (John Wiley & Sons, New York, 1985), "Comprehensive Organic Synthesis. Selectivity, Strategy & Efficiency in Modern Organic Chemistry. In 9 Volumes", Barry M. Trost, Editor-in-Chief (Pergamon Press, New York, 1993 printing).

Generally, the reaction conditions such as temperature, reaction time, solvents, workup procedures, and the like, will be those common in the art for the particular reaction to be performed. The cited reference material, together with material cited therein, contains detailed descriptions of such conditions.

The terms "treated", "treating", "treatment", and the like, mean contacting, mixing, reacting, allowing to react, bringing into contact, and other terms common in the art for indicating that one or more chemical entities is treated in such a manner as to convert it to one or more other chemical entities. This means that "treating compound one with compound two" is synonymous with "allowing compound one to react with compound two", "contacting compound one with compound two", "reacting compound one with compound two", and other expressions common in the art of organic synthesis for reasonably indicating that compound one was "treated", "reacted", "allowed to react", etc., with compound two.

"Treating" indicates the reasonable and usual manner in which organic chemicals are allowed to react. Normal concentrations (0.01M to 10M, typically 0.1M to 1M), temperatures (-100°C to 250°C, typically -78°C to 150°C, more typically -78°C to 100°C, still more typically 0°C to 100°C), solvents (aprotic or protic), reaction times (typically 10 seconds to 10 days, more typically 1 min. to 10 hours, still more typically 10 min. to 6 hours), reaction vessels (typically glass, plastic, metal), pressures, atmospheres

(typically air for oxygen and water insensitive reactions or nitrogen or argon for oxygen or water sensitive), etc., are intended unless otherwise indicated. The knowledge of similar reactions known in the art of organic synthesis are used in selecting the conditions and apparatus for "treating" in a given process. In particular, one of ordinary skill in the art of organic synthesis selects conditions and apparatus reasonably expected to successfully carry out the chemical reactions of the described processes based on the knowledge in the art.


Oxidation and reduction reactions are typically carried out at temperatures near room temperature (about 20°C), although for metal hydride reductions frequently the temperature is reduced to 0°C to -100°C, solvents are typically aprotic for reductions and may be either protic or aprotic for oxidations. Reaction times are adjusted to achieve desired conversions.

Condensation reactions are typically carried out at temperatures near room temperature, although for non-equilibrating, kinetically controlled condensations reduced temperatures (0°C to -100°C) are also common. Solvents can be either protic (common in equilibrating reactions) or aprotic (common in kinetically controlled reactions).

Standard synthetic techniques such as azeotropic removal of reaction by-products and use of anhydrous reaction conditions (e.g. inert gas environments) are common in the art and will be applied when applicable. Workup typically consists of quenching any unreacted reagents followed by partition between a water/organic layer system (extraction) and separating the layer containing the product. Each of the products of the following processes is optionally separated, isolated, and/or purified prior to its use in subsequent processes.

Embodiments

One aspect of the present invention is directed to processes for the preparation of compounds of the formula:




R^1 is a cyclic hydroxy protecting group. A very large number of common protecting groups (including cyclic hydroxy protecting groups) and corresponding chemical cleavage reactions are described in "Protective Groups in Organic Chemistry", Theodora W. Greene (John Wiley & Sons, Inc., New York, 1991, ISBN 0-471-62301-6) ("Greene"). See also Kocienski, Philip J.; "Protecting Groups" (Georg Thieme Verlag Stuttgart, New York, 1994). In particular Chapter 1, Protecting Groups: An Overview, pages 1-20, Chapter 2, Hydroxyl Protecting Groups, pages 21-94, Chapter 3, Diol Protecting Groups, pages 95-117, Chapter 4, Carboxyl Protecting Groups, pages 118-154, Chapter 5, Carbonyl Protecting Groups, pages 155-184, and Chapter 6, Amino Protecting Groups, pages 185-243. Typically, the cyclic hydroxyprotecting groups are those commonly useful as 1,2-diol protecting groups.

Typical 1,2-diol protecting groups (thus, generally where two OH groups are taken together with the R^1 protecting functionality) are described in Greene at pages 118-142 and include Cyclic Acetals and Ketals (Methylene, Ethylidene, 1-*t*-Butylethylidene, 1-Phenylethylidene, (4-Methoxyphenyl)ethylidene, 2,2,2-Trichloroethylidene, Acetonide (Isopropylidene), Cyclopentylidene, Cyclohexylidene, Cycloheptylidene, Benzylidene, *p*-Methoxybenzylidene, 2,4-Dimethoxybenzylidene, 3,4-Dimethoxybenzylidene, 2-Nitrobenzylidene); Cyclic Ortho Esters (Methoxymethylene, Ethoxymethylene, Dimethoxymethylene, 1-Methoxyethylidene, 1-Ethoxyethylidene, 1,2-Dimethoxyethylidene, α -Methoxybenzylidene, 1-(*N,N*-Dimethylamino)ethylidene Derivative, α -(*N,N*-Dimethylamino)benzylidene Derivative, 2-Oxacyclopentylidene); Silyl Derivatives (Di-*t*-butylsilylene Group, 1,3-(1,1,3,3-Tetraisopropylidisiloxanylidene), and Tetra-*t*-butoxydisiloxane-1,3-diylidene), Cyclic Carbonates, Cyclic Boronates, Ethyl Boronate and Phenyl Boronate.

More typically, 1,2-diol protecting groups include those shown in Table A, or alyc ketals or acetals. Still more typically, cyclic ketals and acetals.

5

Table A



wherein R^{1a} is C_1 - C_6 alkyl (as defined immediately below).

"Alkyl" as used herein, unless stated to the contrary, is C_1 - C_6

hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms.

- 10 Examples are methyl (Me, $-\text{CH}_3$), ethyl (Et, $-\text{CH}_2\text{CH}_3$), 1-propyl (n -Pr, n -propyl, $-\text{CH}_2\text{CH}_2\text{CH}_3$), 2-propyl (i -Pr, i -propyl, $-\text{CH}(\text{CH}_3)_2$), 1-butyl (n -Bu, n -butyl, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2-methyl-1-propyl (i -Bu, i -butyl, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$), 2-butyl (s -Bu, s -butyl, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$), 2-methyl-2-propyl (t -Bu, t -butyl, $-\text{C}(\text{CH}_3)_3$), 1-pentyl (n -pentyl, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$),
- 15 2-pentyl ($-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$), 3-pentyl ($-\text{CH}(\text{CH}_2\text{CH}_3)_2$), 2-methyl-2-butyl ($-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_3$), 3-methyl-2-butyl ($-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)_2$), 3-methyl-1-butyl ($-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$), 2-methyl-1-butyl ($-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$), 1-hexyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2-hexyl ($-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3-hexyl ($-\text{CH}(\text{CH}_2\text{CH}_3)(\text{CH}_2\text{CH}_2\text{CH}_3)$),
- 20 2-methyl-2-pentyl ($-\text{C}(\text{CH}_3)_2\text{CH}_2\text{CH}_2\text{CH}_3$), 3-methyl-2-pentyl ($-\text{CH}(\text{CH}_3)\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$), 4-methyl-2-pentyl ($-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}(\text{CH}_3)_2$), 3-methyl-3-pentyl ($-\text{C}(\text{CH}_3)(\text{CH}_2\text{CH}_3)_2$), 2-methyl-3-pentyl ($-\text{CH}(\text{CH}_2\text{CH}_3)\text{CH}(\text{CH}_3)_2$), 2,3-dimethyl-2-butyl ($-\text{C}(\text{CH}_3)_2\text{CH}(\text{CH}_3)_2$), 3,3-dimethyl-2-butyl ($-\text{CH}(\text{CH}_3)\text{C}(\text{CH}_3)_3$). Typical alkyls are methyl, ethyl,
- 25 1-propyl, and 2-propyl.

R^2 is a carboxylic acid protecting group. Typical carboxylic acid protecting groups are R^{25} (described immediately below) or those described in Greene at pages 224-276. Those described in Greene include Esters

- (Methyl); Substituted Methyl Esters (9-Fluorenylmethyl, Methoxymethyl, Methylthiomethyl, Tetrahydropyranyl, Tetrahydrofuranyl, Methoxyethoxymethyl, 2-(Trimethylsilyl)ethoxymethyl, Benzyloxymethyl, Phenacyl, *p*-Bromophenacyl, α -Methylphenacyl, *p*-Methoxyphenacyl, Carboxamidomethyl, *N*-Phthalimidomethyl); 2-Substituted Ethyl Esters (2,2,2-Trichloroethyl, 2-Haloethyl, ω -Chloroalkyl, 2-(Trimethylsilyl)ethyl, 2-Methylthioethyl, 1,3-Dithianyl-2-methyl, 2-(*p*-Nitrophenylsulfenyl)ethyl, 2-(*p*-Toluenesulfonyl)ethyl, 2-(2'-Pyridyl)ethyl, 2-(Diphenylphosphino)ethyl, 1-Methyl-1-phenylethyl, *t*-Butyl, Cyclopentyl, Cyclohexyl, Allyl, 3-Buten-1-yl, 4-(Trimethylsilyl)-2-buten-1-yl, Cinnamyl, α -Methylcinnamyl, Phenyl, *p*-(Methylmercapto)phenyl, Benzyl); Substituted Benzyl Esters (Triphenylmethyl, Diphenylmethyl, Bis(*o*-nitrophenyl)methyl, 9-Anthrylmethyl, 2-(9,10-Dioxo)anthrylmethyl, 5-Dibenzosuberyl, 1-Pyrenylmethyl, 2-(Trifluoromethyl)-6-chromylmethyl, 2,4,6-Trimethylbenzyl, *p*-Bromobenzyl, *o*-Nitrobenzyl, *p*-Nitrobenzyl, *p*-Methoxybenzyl, 2,6-Dimethoxybenzyl, 4-(Methylsulfinyl)benzyl, 4-Sulfobenzyl, Piperonyl, 4-Picolyl, *p*-poly-Benzyl); Silyl Esters (Trimethylsilyl, Triethylsilyl, *t*-Butyldimethylsilyl, *i*-Propyldimethylsilyl, Phenyldimethylsilyl, Di-*t*-butylmethylsilyl); Activated Esters (Thiols); Miscellaneous Derivatives (Oxazoles, 2-Alkyl-1,3-oxazolines, 4-Alkyl-5-oxo-1,3-oxazolidines, 5-Alkyl-4-oxo-1,3-dioxolanes, Ortho Esters, Phenyl Group, Pentaaminocobalt(III) Complex); Stannyl Esters (Triethylstannyl, Tri-*n*-butylstannyl); Amides (*N,N*-Dimethyl, Pyrrolidinyl, Piperidinyl, 5,6-Dihydrophenanthridinyl, *o*-Nitroanilides, *N*-7-Nitroindolyl, *N*-8-Nitro-1,2,3,4-tetrahydroquinolyl, *p*-poly-Benzenesulfonamides); and Hydrazides (Hydrazides, *N*-Phenyl, *N,N'*-Diisopropyl).

R^{25} is alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12 carbon atoms, or alkynyl of 2 to 12 carbon atoms, any one of which alkyl, alkenyl, or alkynyl is substituted with 0-3 R^{22} groups (R^{22} is described below). More typically R^{25} is alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, or alkynyl of 2 to 6 carbon atoms, any one of which alkyl, alkenyl, or alkynyl is substituted with 0-3 R^{22} groups. Still more typically, R^{25} is alkyl of 1 to 8 carbon atoms substituted with 0-2 R^{22} groups. Even more typically, R^{25} is alkyl of 1, 2, 3, 4, 5, 6, 7, or 8 carbon atoms. Most typically R^{25} is methyl, ethyl, 1-propyl or 2-propyl.

"Alkenyl" as used herein, unless stated to the contrary, is C₁-C₆ hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms.

Examples are ethenyl ($-\text{CH}=\text{CH}_2$), 1-prop-1-enyl ($-\text{CH}=\text{CHCH}_3$), 1-prop-2-enyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), 2-prop-1-enyl ($-\text{C}(=\text{CH}_2)(\text{CH}_3)$), 1-but-1-enyl ($-\text{CH}=\text{CHCH}_2\text{CH}_3$), 1-but-2-enyl ($-\text{CH}_2\text{CH}=\text{CHCH}_3$), 1-but-3-enyl ($-\text{CH}_2\text{CH}_2\text{CH}=\text{CH}_2$), 2-methyl-1-prop-1-enyl ($-\text{CH}=\text{C}(\text{CH}_3)_2$), 2-methyl-1-prop-2-enyl ($-\text{CH}_2\text{C}(=\text{CH}_2)(\text{CH}_3)$), 2-but-1-enyl ($-\text{C}(=\text{CH}_2)\text{CH}_2\text{CH}_3$), 2-but-2-enyl ($-\text{C}(\text{CH}_3)=\text{CHCH}_3$), 2-but-3-enyl ($-\text{CH}(\text{CH}_3)\text{CH}=\text{CH}_2$), 1-pent-1-enyl ($-\text{C}=\text{CHCH}_2\text{CH}_2\text{CH}_3$), 1-pent-2-enyl ($-\text{CHCH}=\text{CHCH}_2\text{CH}_3$), 1-pent-3-enyl ($-\text{CHCH}_2\text{CH}=\text{CHCH}_3$), 1-pent-4-enyl ($-\text{CHCH}_2\text{CH}_2\text{CH}=\text{CH}_2$), 2-pent-1-enyl ($-\text{C}(=\text{CH}_2)\text{CH}_2\text{CH}_2\text{CH}_3$), 2-pent-2-enyl ($-\text{C}(\text{CH}_3)=\text{CH}_2\text{CH}_2\text{CH}_3$), 2-pent-3-enyl ($-\text{CH}(\text{CH}_3)\text{CH}=\text{CHCH}_3$), 2-pent-4-enyl ($-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}=\text{CH}_2$) or 3-methyl-1-but-2-enyl ($-\text{CH}_2\text{CH}=\text{C}(\text{CH}_3)_2$). More typically, alkenyl groups are of 2, 3 or 4 carbon atoms.

"Alkynyl" as used herein, unless stated to the contrary, is C_1 - C_6 hydrocarbon containing normal, secondary, tertiary or cyclic carbon atoms. Examples are ethynyl ($-\text{C}\equiv\text{CH}$), 1-prop-1-ynyl ($-\text{C}\equiv\text{CCH}_3$), 1-prop-2-ynyl ($-\text{CH}_2\text{C}\equiv\text{CH}$), 1-but-1-ynyl ($-\text{C}\equiv\text{CCH}_2\text{CH}_3$), 1-but-2-ynyl ($-\text{CH}_2\text{C}\equiv\text{CCH}_3$), 1-but-3-ynyl ($-\text{CH}_2\text{CH}_2\text{C}\equiv\text{CH}$), 2-but-3-ynyl ($\text{CH}(\text{CH}_3)\text{C}\equiv\text{CH}$), 1-pent-1-ynyl ($-\text{C}\equiv\text{CCH}_2\text{CH}_2\text{CH}_3$), 1-pent-2-ynyl ($-\text{CH}_2\text{C}\equiv\text{CCH}_2\text{CH}_3$), 1-pent-3-ynyl ($-\text{CH}_2\text{CH}_2\text{C}\equiv\text{CCH}_3$) or 1-pent-4-ynyl ($-\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CH}$). More typically, alkynyl groups are of 2, 3 or 4 carbon atoms.

R^3 is a hydroxy protecting group. Typical R^3 hydroxy protecting groups described in Greene (pages 14-118) include Ethers (Methyl); Substituted Methyl Ethers (Methoxymethyl, Methylthiomethyl, *t*-Butylthiomethyl, (Phenyldimethylsilyl)methoxymethyl, Benzyloxymethyl, *p*-Methoxybenzyloxymethyl, (4-Methoxyphenoxy)methyl, Guaiacolmethyl, *t*-Butoxymethyl, 4-Pentenylloxymethyl, Siloxymethyl, 2-Methoxyethoxymethyl, 2,2,2-Trichloroethoxymethyl, Bis(2-chloroethoxy)methyl, 2-(Trimethylsilyl)ethoxymethyl, Tetrahydropyranyl, 3-Bromotetrahydropyranyl, Tetrahydrothiopyranyl, 1-Methoxycyclohexyl, 4-Methoxytetrahydropyranyl, 4-Methoxytetrahydrothiopyranyl, 4-Methoxytetrahydrothiopyranyl *S,S*-Dioxido, 1-[(2-Chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl, 3,5,1,4-Dioxan-2-yl, Tetrahydrofuran-2-yl, Tetrahydrothiofuran-2-yl, 2,3,3a,4,5,6,7,7a-Octahydro-7,8,8-trimethyl-4,7-methanobenzofuran-2-yl); Substituted Ethyl Ethers (1-Ethoxyethyl, 1-(2-Chloroethoxy)ethyl, 1-Methyl-1-methoxyethyl, 1-Methyl-1-benzyloxyethyl, 1-Methyl-1-benzyloxy-2-fluoroethyl, 2,2,2-Trichloroethyl,


- 2-Trimethylsilylethyl, 2-(Phenylselenyl)ethyl, *t*-Butyl, Allyl, *p*-Chlorophenyl, *p*-Methoxyphenyl, 2,4-Dinitrophenyl, Benzyl); Substituted Benzyl Ethers (*p*-Methoxybenzyl, 3,4-Dimethoxybenzyl, *o*-Nitrobenzyl, *p*-Nitrobenzyl, *p*-Halobenzyl, 2,6-Dichlorobenzyl, *p*-Cyanobenzyl, *p*-Phenylbenzyl, 2- and 4-Picolyl, 3-Methyl-2-picolyl *N*-Oxido, 5 Diphenylmethyl, *p,p'*-Dinitrobenzhydryl, 5-Dibenzosuberyl, Triphenylmethyl, α -Naphthyldiphenylmethyl, *p*-methoxyphenyldiphenylmethyl, Di(*p*-methoxyphenyl)phenylmethyl, Tri(*p*-methoxyphenyl)methyl, 4-(4'-Bromophenacyloxy)phenyldiphenylmethyl, 4,4',4''-Tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-Tris(levulinoyloxyphenyl)methyl, 4,4',4''-Tris(benzoyloxyphenyl)methyl, 3-(Imidazol-1-ylmethyl)bis(4',4''-dimethoxyphenyl)methyl, 1,1-Bis(4-methoxyphenyl)-1'-pyrenylmethyl, 9-Anthryl, 9-(9-Phenyl)xanthenyl, 9-(9-Phenyl-10-oxo)anthryl, 1,3-Benzodithiolan-2-yl, Benzisothiazolyl *S,S*-Dioxido); Silyl Ethers (Trimethylsilyl, Triethylsilyl, Triisopropylsilyl, Dimethylisopropylsilyl, Diethylisopropylsilyl, Dimethylthexylsilyl, *t*-Butyldimethylsilyl, *t*-Butyldiphenylsilyl, Tribenzylsilyl, Tri-*p*-xylylsilyl, Triphenylsilyl, Diphenylmethylsilyl, *t*-Butylmethoxyphenylsilyl); Esters (Formate, Benzoylformate, Acetate, Chloroacetate, Dichloroacetate, Trichloroacetate, Trifluoroacetate, Methoxyacetate, Triphenylmethoxyacetate, Phenoxyacetate, *p*-Chlorophenoxyacetate, *p*-poly-Phenylacetate, 3-Phenylpropionate, 4-Oxopentanoate (Levulinate), 4,4-(Ethylenedithio)pentanoate, Pivaloate, Adamantoate, Crotonate, 4-Methoxycrotonate, Benzoate, *p*-Phenylbenzoate, 2,4,6-Trimethylbenzoate (Mesitoate)); Carbonates (Methyl, 9-Fluorenylmethyl, Ethyl, 2,2,2-Trichloroethyl, 2-(Trimethylsilyl)ethyl, 2-(Phenylsulfonyl)ethyl, 2-(Triphenylphosphonio)ethyl, Isobutyl, Vinyl, Allyl, *p*-Nitrophenyl, Benzyl, *p*-Methoxybenzyl, 3,4-Dimethoxybenzyl, *o*-Nitrobenzyl, *p*-Nitrobenzyl, *S*-Benzyl Thiocarbonate, 4-Ethoxy-1-naphthyl, Methyl Dithiocarbonate); Groups With Assisted Cleavage (2-Iodobenzoate, 4-Azidobutyrate, 4-Nitro-4-methylpentanoate, *o*-(Dibromomethyl)benzoate, 2-Formylbenzenesulfonate, 2-(Methylthiomethoxy)ethyl Carbonate, 4-(Methylthiomethoxy)butyrate, 2-(Methylthiomethoxymethyl)benzoate); Miscellaneous Esters (2,6-Dichloro-4-methylphenoxyacetate, 2,6-Dichloro-4-(1,1,3,3-tetramethylbutyl)phenoxyacetate, 2,4-Bis(1,1-

dimethylpropyl)phenoxyacetate, Chorodiphenylacetate, Isobutyrate, Monosuccinoate, (E)-2-Methyl-2-butenate (Tigloate), *o*-(Methoxycarbonyl)benzoate, *p*-poly-Benzoate, α -Naphthoate, Nitrate, Alkyl *N,N,N',N'*-Tetramethylphosphorodiamidate, *N*-Phenylcarbamate, Borate, Dimethylphosphinothioyl, 2,4-Dinitrophenylsulfenate); and Sulfonates (Sulfate, Methanesulfonate (Mesylate), Benzylsulfonate, Tosylate).

More typically, R^3 hydroxy protecting groups include substituted methyl ethers, substituted benzyl ethers, silyl ethers, and esters including sulfonic acid esters, still more typically, trialkylsilyl ethers, tosylates, mesylates and acetates.

Each R^{20} is independently H or an alkyl of 1 to 12 carbon atoms. Typically R^{20} is H or alkyl of 1 to 6 carbon as described above. Still more typically, R^{20} is H or methyl. More typically yet, R^{20} is H.

This process embodiment comprises reaction of a compound of the formula:



with a dehydrating reagent. Typically the hydroxy group at position 1 is eliminated without removing the cis-4,5-diol protecting group. The hydroxy group at position 1 is eliminated to form an olefinic bond between positions 1 and 6.

Typically the process comprises treating compound 4 with a suitable dehydrating agent, such as a mineral acid (HCl, H₂SO₄) or SO₂Cl₂. More typically, compound 4 is treated with SO₂Cl₂, followed by an alkanol. Still more typically, compound 4 is treated with SO₂Cl₂ in a suitable polar, aprotic solvent, such as an amine to form an olefin. More typically yet, compound 4 is treated with SO₂Cl₂ in pyridine/CH₂Cl₂ at a temperature between -100°C and 0°C, typically -100°C and -10°C, more typically -78°C, to form compound 5.


In a typical embodiment, a solution of compound 4 and pyridine in dichloromethane is cooled to -20° to -30°C and treated portionwise with

sulfuryl chloride. After the exothermic reaction subsided, the resulting slurry is quenched with ethanol, warmed to 0°C, and washed successively with 16% sulfuric acid, water and 5% aqueous sodium bicarbonate. A detailed example of this embodiment is provided as Example 4 below.

5 Optionally, the process of this embodiment further comprises purifying or separating compound 5 from any other reaction products or other contaminants such as other double bond isomers, halogenated side products or starting materials and reagents by treatment with a noble metal complex. Noble metals include gold, silver, platinum, palladium, 10 iridium, rhenium, mercury, ruthenium and osmium. Typically, the noble metal complex of this embodiment is a complex of platinum or palladium. More typically the complex is a palladium (0) complex, still more typically, the complex is a tetrakis(triarylphosphine)palladium (0) complex.

15 In a typical embodiment the organic layer of the reaction contains a mixture of olefin and halogenated products as well as starting material. It is concentrated *in vacuo* and ethyl acetate is added. The solution is treated with pyrrolidine and tetrakis(triphenylphosphine)palladium(0) at ambient temperature, followed by washing with 16% sulfuric acid. The organic layer is filtered through a pad of silica gel and eluted with ethyl acetate. 20 The filtrate is concentrated *in vacuo*. The residue is dissolved in ethyl acetate at reflux and hexane is added. Upon cooling, the product crystallizes and is separated by filtration and washed with 14% ethyl acetate in hexane. After drying *in vacuo*, 5 was obtained. A detailed example of this embodiment is provided as Example 4 below.


25 In another example of this embodiment compound 5 is of the formula:



6

Another aspect of the present invention is directed to processes for the preparation of compounds of the formula:

30



11

wherein:

R^2 , R^3 and R^{20} are as defined above.

R^4 is described below.

- 5 W is carbocycle or heterocycle wherein any one of which carbocycle or heterocycle is substituted with 0 to 3 R^{29} groups (R^{29} is described below).

W is a carbocycle or heterocycle, with the proviso that each W is independently substituted with 0 to 3 R^{29} groups (R^{29} is described below). W carbocycles and heterocycles are stable chemical structures. Such
 10 structures are isolatable in measurable yield, with measurable purity, from reaction mixtures at temperatures from -78°C to 200°C . Each W is independently substituted with 0 to 3 R^{29} groups. Typically, W is a saturated, unsaturated or aromatic ring comprising a mono- or bicyclic carbocycle or heterocycle. More typically, W has 3 to 10 ring atoms, still
 15 more typically, 3 to 7 ring atoms, and ordinarily 3 to 6 ring atoms. The W rings are saturated when containing 3 ring atoms, saturated or monounsaturated when containing 4 ring atoms, saturated, or mono- or diunsaturated when containing 5 ring atoms, and saturated, mono- or diunsaturated, or aromatic when containing 6 ring atoms.

- 20 When W is carbocyclic, it is typically a 3 to 7 carbon monocycle or a 7 to 12 carbon atom bicycle. More typically, W monocyclic carbocycles have 3 to 6 ring atoms, still more typically 5 or 6 ring atoms. W bicyclic carbocycles typically have 7 to 12 ring atoms arranged as a bicyclo [4,5], [5,5], [5,6] or [6,6] system, still more typically, 9 or 10 ring atoms arranged as a
 25 bicyclo [5,6] or [6,6] system. Examples include cyclopropyl, cyclobutyl, cyclopentyl, 1-cyclopent-1-enyl, 1-cyclopent-2-enyl, 1-cyclopent-3-enyl, cyclohexyl, 1-cyclohex-1-enyl, 1-cyclohex-2-enyl, 1-cyclohex-3-enyl, phenyl, spiryl and naphthyl.

- 30 A W heterocycle is typically a monocycle having 3 to 7 ring members (2 to 6 carbon atoms and 1 to 3 heteroatoms selected from N, O, P, and S) or a bicycle having 7 to 10 ring members (4 to 9 carbon atoms and

1 to 3 heteroatoms selected from N, O, P, and S). More typically, W heterocyclic monocycles have 3 to 6 ring atoms (2 to 5 carbon atoms and 1 to 2 heteroatoms selected from N, O, and S), still more typically, 5 or 6 ring atoms (3 to 5 carbon atoms and 1 to 2 heteroatoms selected from N and S).

5 W heterocyclic bicycles typically have 7 to 10 ring atoms (6 to 9 carbon atoms and 1 to 2 heteroatoms selected from N, O, and S) arranged as a bicyclo [4,5], [5,5], [5,6], or [6,6] system, still more typically, 9 to 10 ring atoms (8 to 9 carbon atoms and 1 to 2 hetero atoms selected from N and S) arranged as a bicyclo [5,6] or [6,6] system.

10 "Heterocycle" as used herein includes by way of example and not limitation these heterocycles described in Paquette, Leo A.; "Principles of Modern Heterocyclic Chemistry" (W.A. Benjamin, New York, 1968), particularly Chapters 1, 3, 4, 6, 7, and 9; "The Chemistry of Heterocyclic Compounds, A series of Monographs" (John Wiley & Sons, New York,
15 1950 to present), in particular Volumes 13, 14, 16, 19, and 28; and "J. Am. Chem. Soc.", 82:5566 (1960).

Examples of heterocycles include by way of example and not limitation pyridyl, thiazolyl, tetrahydrothiophenyl, sulfur oxidized tetrahydrothiophenyl, pyrimidinyl, furanyl, thienyl, pyrrolyl, pyrazolyl,
20 imidazolyl, tetrazolyl, benzofuranyl, thianaphthalenyl, indolyl, indolenyl, quinoliny, isoquinoliny, benzimidazolyl, piperidinyl, 4-piperidonyl, pyrrolidinyl, 2-pyrrolidonyl, pyrrolinyl, tetrahydrofuranyl, tetrahydroquinoliny, tetrahydroisoquinoliny, decahydroquinoliny,
25 octahydroisoquinoliny, azociny, triazinyl, 6H-1,2,5-thiadiazinyl, 2H,6H-1,5,2-dithiazinyl, thienyl, thianthrenyl, pyranyl, isobenzofuranyl, chromenyl, xanthenyl, phenoxathiinyl, 2H-pyrrolyl, isothiazolyl, isoxazolyl, pyrazinyl, pyridazinyl, indoliziny, isoindolyl, 3H-indolyl, 1H-indazoly, purinyl, 4H-quinoliziny, phthalazinyl, naphthyridinyl, quinoxaliny, quinazoliny, cinnoliny, pteridinyl, 4aH-carbazolyl,
30 carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, pyrimidinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, furazanyl, phenoxazinyl, isochromanyl, chromanyl, imidazolidinyl, imidazoliny, pyrazolidinyl, pyrazoliny, piperazinyl, indoliny, isoindoliny, quinuclidiny, morpholiny, oxazolidinyl, benzotriazolyl, benzisoxazolyl, oxindolyl,
35 benzoxazoliny, and isatinoyl.

By way of example and not limitation, carbon bonded heterocycles are bonded at position 2, 3, 4, 5, or 6 of a pyridine, position 3, 4, 5, or 6 of a


pyridazine, position 2, 4, 5, or 6 of a pyrimidine, position 2, 3, 5, or 6 of a pyrazine, position 2, 3, 4, or 5 of a furan, tetrahydrofuran, thiofuran, thiophene, pyrrole or tetrahydropyrrole, position 2, 4, or 5 of an oxazole, imidazole or thiazole, position 3, 4, or 5 of an isoxazole, pyrazole, or
5 isothiazole, position 2 or 3 of an aziridine, position 2, 3, or 4 of an azetidine, position 2, 3, 4, 5, 6, 7, or 8 of a quinoline or position 1, 3, 4, 5, 6, 7, or 8 of an isoquinoline. Still more typically, carbon bonded heterocycles include 2-pyridyl, 3-pyridyl, 4-pyridyl, 5-pyridyl, 6-pyridyl, 3-pyridazinyl, 4-pyridazinyl, 5-pyridazinyl, 6-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl, 2-pyrazinyl, 3-pyrazinyl, 5-pyrazinyl, 6-pyrazinyl, 2-thiazolyl, 4-thiazolyl, or 5-thiazolyl.
10

By way of example and not limitation, nitrogen bonded heterocycles are bonded at position 1 of an aziridine, azetidine, pyrrole, pyrrolidine, 2-pyrroline, 3-pyrroline, imidazole, imidazolidine, 2-imidazoline, 3-imidazoline, pyrazole, pyrazoline, 2-pyrazoline, 3-pyrazoline, piperidine,
15 piperazine, indole, indoline, 1H-indazole, position 2 of a isoindole, or isoindoline, position 4 of a morpholine, and position 9 of a carbazole, or β -carboline. Still more typically, nitrogen bonded heterocycles include 1-aziridyl, 1-azetetyl, 1-pyrrolyl, 1-imidazolyl, 1-pyrazolyl, and 1-piperidinyl.
20

Typically W heterocycles are selected from pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, s-triazinyl, oxazolyl, imidazolyl, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, furanyl, thiofuranyl, thienyl, or pyrrolyl.

More typically, the heterocycle of W is bonded through a carbon atom or nitrogen atom thereof. Still more typically W heterocycles are
25 bonded by a stable covalent bond through a carbon or nitrogen atom thereof. Stable covalent bonds are chemically stable structures as described above.

W optionally is selected from the group consisting of:



R⁵ is H or R³.

- R⁷ is H or an amino protecting group. R⁷ amino protecting groups are described by Greene at pages 315-385. They include Carbamates (methyl and ethyl, 9-fluorenylmethyl, 9(2-sulfo)fluorenylmethyl, 9-(2,7-dibromo)fluorenylmethyl, 2,7-di-*t*-buthyl-[9-(10,10-dioxo-10,10,10,10-tetrahydrothioxanthyl)]methyl, 4-methoxyphenacyl); Substituted Ethyl (2,2,2-trichloroethyl, 2-trimethylsilyl ethyl, 2-phenylethyl, 1-(1-adamantyl)-1-methylethyl, 1,1-dimethyl-2-haloethyl, 1,1-dimethyl-2,2-dibromoethyl, 1,1-dimethyl-2,2,2-trichloroethyl, 1-methyl-1-(4-biphenyl)ethyl, 1-(3,5-di-*t*-butylphenyl)-1-methylethyl, 2-(2'- and 4'-pyridyl)ethyl, 2-(*N,N*-dicyclohexylcarboxamido)ethyl, *t*-butyl, 1-adamantyl, vinyl, allyl, 1-isopropylallyl, cinnamyl, 4-nitrocinnamyl, 8-quinolyl, *N*-hydroxypiperidinyl, alkyl dithio, benzyl, *p*-methoxybenzyl, *p*-nitrobenzyl, *p*-bromobenzyl, *p*-chlorobenzyl, 2,4-dichlorobenzyl, 4-methylsulfinylbenzyl, 9-anthrylmethyl, diphenylmethyl); Groups With Assisted Cleavage (2-methylthioethyl, 2-methylsulfonyl ethyl, 2-(*p*-toluenesulfonyl)ethyl, [2-(1,3-dithianyl)]methyl, 4-methylthiophenyl, 2,4-dimethylthiophenyl, 2-phosphonioethyl, 2-triphenylphosphonioisopropyl, 1,1-dimethyl-2-cyanoethyl, *m*-chloro-*p*-acyloxybenzyl, *p*-(dihydroxyboryl)benzyl, 5-benzisoxazolylmethyl, 2-(trifluoromethyl)-6-chromonylmethyl); Groups Capable of Photolytic Cleavage (*m*-nitrophenyl, 3,5-dimethoxybenzyl, *o*-nitrobenzyl, 3,4-dimethoxy-6-nitrobenzyl, phenyl(*o*-nitrophenyl)methyl); Urea-Type Derivatives (phenothiazinyl-(10)-carbonyl, *N'*-*p*-toluenesulfonylamino carbonyl, *N'*-phenylaminothiocarbonyl);

Miscellaneous Carbamates (*t*-amyl, *S*-benzyl thiocarbamate, *p*-cyanobenzyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopropylmethyl, *p*-decyloxybenzyl, diisopropylmethyl, 2,2-dimethoxycarbonylvinyl, *o*-(*N,N*-dimethylcarboxamido)benzyl, 1,1-dimethyl-3-(*N,N*-dimethylcarboxamido)propyl, 1,1-dimethylpropynyl, di(2-pyridyl)methyl, 2-furanylmethyl, 2-Iodoethyl, Isobornyl, Isobutyl, Isonicotinyl, *p*-(*p*'-Methoxyphenylazo)benzyl, 1-methylcyclobutyl, 1-methylcyclohexyl, 1-methyl-1-cyclopropylmethyl, 1-methyl-1-(3,5-dimethoxyphenyl)ethyl, 1-methyl-1-(*p*-phenylazophenyl)ethyl, 1-methyl-1-phenylethyl, 1-methyl-1-(4-pyridyl)ethyl, phenyl, *p*-(phenylazo)benzyl, 2,4,6-tri-*t*-butylphenyl, 4-(trimethylammonium)benzyl, 2,4,6-trimethylbenzyl); Amides (*N*-formyl, *N*-acetyl, *N*-choroacetyl, *N*-trichoroacetyl, *N*-trifluoroacetyl, *N*-phenylacetyl, *N*-3-phenylpropionyl, *N*-picolinoyl, *N*-3-pyridylcarboxamide, *N*-benzoylphenylalanyl, *N*-benzoyl, *N*-*p*-phenylbenzoyl); Amides With Assisted Cleavage (*N*-*o*-nitrophenylacetyl, *N*-*o*-nitrophenoxycetyl, *N*-acetoacetyl, (*N*'-dithiobenzoyloxycarbonylamino)acetyl, *N*-3-(*p*-hydroxyphenyl)propionyl, *N*-3-(*o*-nitrophenyl)propionyl, *N*-2-methyl-2-(*o*-nitrophenoxy)propionyl, *N*-2-methyl-2-(*o*-phenylazophenoxy)propionyl, *N*-4-chlorobutyryl, *N*-3-methyl-3-nitrobutyryl, *N*-*o*-nitrocinnamoyl, *N*-acetylmethionine, *N*-*o*-nitrobenzoyl, *N*-*o*-(benzoyloxymethyl)benzoyl, 4,5-diphenyl-3-oxazolin-2-one); Cyclic Imide Derivatives (*N*-phthalimide, *N*-dithiasuccinoyl, *N*-2,3-diphenylmaleoyl, *N*-2,5-dimethylpyrrolyl, *N*-1,1,4,4-tetramethyldisilylazacyclopentane adduct, 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, 1-substituted 3,5-dinitro-4-pyridonyl); *N*-Alkyl and *N*-Aryl Amines (*N*-methyl, *N*-allyl, *N*-[2-(trimethylsilyl)ethoxy]methyl, *N*-3-acetoxypentyl, *N*-(1-isopropyl-4-nitro-2-oxo-3-pyrrolin-3-yl), Quaternary Ammonium Salts, *N*-benzyl, *N*-di(4-methoxyphenyl)methyl, *N*-5-dibenzosuberyl, *N*-triphenylmethyl, *N*-(4-methoxyphenyl)diphenylmethyl, *N*-9-phenylfluorenyl, *N*-2,7-dichloro-9-fluorenylmethylene, *N*-ferrocenylmethyl, *N*-2-picolylamine *N*'-oxide), Imine Derivatives (*N*-1,1-dimethylthiomethylene, *N*-benzylidene, *N*-*p*-methoxybenzylidene, *N*-diphenylmethylene, *N*-[(2-pyridyl)mesityl]methylene, *N*-(*N*',*N*'-dimethylamino)methylene, *N*,*N*'-isopropylidene, *N*-*p*-nitrobenzylidene, *N*-salicylidene, *N*-5-chlorosalicylidene, *N*-(5-chloro-2-hydroxyphenyl)phenylmethylene, *N*-cyclohexylidene); Enamine Derivatives (*N*-(5,5-dimethyl-3-oxo-1-cyclohexenyl)); *N*-Metal Derivatives

(*N*-borane derivatives, *N*-diphenylborinic acid derivatives, *N*-[phenyl(pentacarbonylchromium- or -tungsten)]carbenyl, *N*-copper or *N*-zinc chelate); *N*-*N* Derivatives (*N*-nitro, *N*-nitroso, *N*-oxide); *N*-*P* Derivatives (*N*-diphenylphosphinyl, *N*-dimethylthiophosphinyl, *N*-diphenylthiophosphinyl, *N*-dialkyl phosphoryl, *N*-dibenzyl phosphoryl, *N*-diphenyl phosphoryl); *N*-*Si* Derivatives; *N*-*S* Derivatives; *N*-Sulfenyl Derivatives (*N*-benzenesulfenyl, *N*-*o*-nitrobenzenesulfenyl, *N*-2,4-dinitrobenzenesulfenyl, *N*-pentachlorobenzenesulfenyl, *N*-2-nitro-4-methoxybenzenesulfenyl, *N*-triphenylmethylsulfenyl, *N*-3-nitropyridinesulfenyl); and *N*-sulfonyl Derivatives (*N*-*p*-toluenesulfonyl, *N*-benzenesulfonyl, *N*-2,3,6-trimethyl-4-methoxybenzenesulfonyl, *N*-2,4,6-trimethoxybenzenesulfonyl, *N*-2,6-dimethyl-4-methoxybenzenesulfonyl, *N*-pentamethylbenzenesulfonyl, *N*-2,3,5,6-tetramethyl-4-methoxybenzenesulfonyl, *N*-4-methoxybenzenesulfonyl, *N*-2,4,6-trimethylbenzenesulfonyl, *N*-2,6-dimethoxy-4-methylbenzenesulfonyl, *N*-2,2,5,7,8-pentamethylchroman-6-sulfonyl, *N*-methanesulfonyl, *N*- β -trimethylsilyethanesulfonyl, *N*-9-anthracenesulfonyl, *N*-4-(4',8'-dimethoxynaphthylmethyl)benzenesulfonyl, *N*-benzylsulfonyl, *N*-trifluoromethylsulfonyl, *N*-phenacylsulfonyl). Typically, R^7 is H or a -C(O) R^{25} (R^{25} is described above).

R^8 is H or R^2 . Typically R^8 is H.

R^9 is H or a thiol protecting group. R^9 amino protecting groups are described by Greene at pages 277-308. They include Thioethers (*S*-Benzyl, *S*-*p*-Methoxybenzyl, *S*-*o*- or *p*-Hydroxy- or Acetoxybenzyl, *S*-*p*-Nitrobenzyl, *S*-4-Picolyl, *S*-2-Picolyl *N*-Oxide, *S*-9-Anthrylmethyl, *S*-9-Fluorenylmethyl, *S*-Ferrocenylmethyl); *S*-Diphenylmethyl, Substituted *S*-Diphenylmethyl, and *S*-Triphenylmethyl Thioethers (*S*-Diphenylmethyl, *S*-Bis(4-methoxyphenyl)methyl, *S*-5-Dibenzosuberyl, *S*-Triphenylmethyl, *S*-Diphenyl-4-pyridylmethyl, *S*-Phenyl, *S*-2,4-Dinitrophenyl, *S*-*t*-Butyl, *S*-1-Adamantyl); Substituted *S*-Methyl Derivatives Monothio, Dithio, and Amino thio Acetals (*S*-Methoxymethyl, *S*-Isobutoxymethyl, *S*-2-Tetrahydropyranyl, *S*-Benzylthiomethyl, *S*-Phenylthiomethyl, Thiazolidines, *S*-Acetamidomethyl, *S*-Trimethylacetamidomethyl, *S*-Benzamidomethyl, *S*-Acetyl-, *S*-Carboxy-, and *S*-Cyanomethyl); Substituted *S*-Ethyl Derivatives (*S*-2-Nitro-1-phenylethyl, *S*-2-(4'-Pyridyl)ethyl, *S*-2-Cyanoethyl, *S*-2,2-Bis(carboethoxy)ethyl, *S*-1-*m*-Nitrophenyl-2-benzoyl ethyl, *S*-2-Phenylsulfonyl ethyl, *S*-1-(4-Methylphenylsulfonyl)-2-

- methylprop-2-yl); Silyl Thioethers, Thioesters, (S-Acetyl Derivative, S-Benzoyl Derivative, S-N-[(p-Biphenyl)isopropoxy]carbonyl]-N-methyl- γ -aminobutyrate, S-N-(t-Butoxycarbonyl)-N-methyl- γ -aminobutyrate); Thiocarbonate Derivatives (S-2,2,2-Trichloroethoxycarbonyl, S-t-Butoxycarbonyl, S-Benzoyloxycarbonyl, S-p-Methoxybenzyloxycarbonyl); Thiocarbamate Derivatives (S-(N-Ethyl), S-(N-Methoxymethyl); Miscellaneous Derivatives, Unsymmetrical Disulfides (S-Ethyl, S-t-Butyl, Substituted S-Phenyl); Sulfenyl Derivatives (S-Sulfonate, S-Sulfenylthiocarbonate, S-3-Nitro-2-pyridinesulfenyl Sulfide); Protection for Dithiols, Dithio Acetals and Ketals (S,S'-Methylene, S,S'-Isopropylidene, and S,S'-Benzylidene, S,S'-p-Methoxybenzylidene); Protection for Sulfides (S-Methylsulfonium Salt, S-Benzyl- and S-4-Methoxybenzylsulfonium Salt, S-1-(4-Phthalimidobutyl)sulfonium Salt); S-P Derivatives (S-(Dimethylphosphino)thioyl, S-(Diphenylphosphino)thioyl);

Each R^{21} is independently R^{20} , Br, Cl, F, I, CN, NO_2 or N_3 . Typically, R^{21} is Cl, F or R^{20} , more typically, R^{20} , still more typically, H.

- Each R^{22} is independently F, Cl, Br, I, -CN, N_3 , - NO_2 , -OR⁵, -OR²⁰, -N(R^{20})₂, -N(R^{20})(R^7), -N(R^7)₂, -SR²⁰, -SR⁹, -S(O) R^{20} , -S(O)₂ R^{20} , -S(O)OR²⁰, -S(O)OR⁸, -S(O)₂OR²⁰, -S(O)₂OR⁸, -C(O)OR²⁰, -C(O)OR⁸, -OC(O) R^{20} , -N(R^{20})(C(O) R^{20}), -N(R^7)(C(O) R^{20}), -N(R^{20})(C(O)OR²⁰), -N(R^7)(C(O)OR²⁰), -C(O)N(R^{20})₂, -C(O)N(R^7)(R^{20}), -C(O)N(R^7)₂, -C(NR²⁰)(N(R^{20})₂), -C(N(R^7))(N(R^{20})₂), -C(N(R^{20}))(N(R^{20})(R^7)), -C(N(R^7))(N(R^{20})(R^7)), -C(N(R^{20}))(N(R^7)₂), -C(N(R^7))(N(R^7)₂), -N(R^{20})C(N(R^{20}))(N(R^{20})₂), -N(R^{20})C(N(R^{20}))(N(R^{20})(R^7)), -N(R^{20})C(N(R^7))(N(R^{20})₂), -N(R^7)C(N(R^{20}))(N(R^{20})₂), -N(R^7)C(N(R^7))(N(R^{20})₂), -N(R^7)C(N(R^{20}))(N(R^{20})(R^7)), -N(R^{20})C(N(R^7))(N(R^{20})(R^7)), -N(R^{20})C(N(R^{20}))(N(R^7)₂), -N(R^7)C(N(R^7))(N(R^{20})(R^7)), -N(R^7)C(N(R^{20}))(N(R^7)₂), -N(R^{20})C(N(R^7))(N(R^7)₂), -N(R^7)C(N(R^7))(N(R^7)₂), =O, =S, =N(R^{20}), =N(R^7) or W.

- Typically R^{22} is F, Cl, Br, I, -CN, N_3 , - NO_2 , -OR⁵, -OR²⁰, -N(R^{20})₂, -N(R^{20})(R^7), -N(R^7)₂, -SR²⁰, -SR⁹, -S(O) R^{20} , -S(O)₂ R^{20} , -S(O)OR²⁰, -S(O)OR⁸, -S(O)₂OR²⁰, -S(O)₂OR⁸, -C(O)OR²⁰, -C(O)OR⁸, =O, =S, =N(R^{20}) or =N(R^7). More typically, R^{22} is F, Cl, Br, -CN, N_3 , - NO_2 , -OR⁵, -OR²⁰, -N(R^{20})₂, -N(R^{20})(R^7), -N(R^7)₂, -C(O)OR²⁰, -C(O)OR⁸, or =O. Still more typically R^{22} is F, Cl, Br, -CN, N_3 , - NO_2 , -OR²⁰, -N(R^{20})₂, -C(O)OR²⁰ or =O. More typically yet R^{22} is F, Cl, Br, -CN, -OH, -N(H)₂, -C(O)OR²⁰ or =O.

Each R^{23} is independently alkyl of 1 to 11 carbon atoms, alkenyl of 2 to 11 carbon atoms, or alkynyl of 2 to 11 carbon atoms. More typically R^{23} is alkyl of 1 to 8 carbon atoms, alkenyl of 2 to 8 carbon atoms, or alkynyl of 2 to 8 carbon atoms, still more typically, R^{23} is alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, or alkynyl of 2 to 6 carbon atoms. More typically yet, R^{23} is R^{25} .

Each R^{24} is independently R^{23} wherein each R^{23} is substituted with 0 to 3 R^{22} groups. Each of the typical embodiments of R^{23} and R^{22} are typical of R^{24} . More typically R^{24} is substituted with 0, 1, 2, or 3 R^{22} groups.

R^{24a} is independently alkylene of 1 to 11 carbon atoms, alkenylene of 2 to 11 carbon atoms, or alkynylene of 2-11 carbon atoms any one of which alkylene, alkenylene or alkynylene is substituted with 0-3 R^{22} groups. More typically R^{24a} is alkylene of 1 to 8 carbon atoms, alkenylene of 2 to 8 carbon atoms, or alkynylene of 2 to 8 carbon atoms, still more typically, R^{24a} is alkylene of 1 to 6 carbon atoms, alkenylene of 2 to 6 carbon atoms, or alkynylene of 2 to 6 carbon atoms. More typically yet, R^{24a} is $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$ or $-C(H)(CH_3)-$.

Each R^{28} is independently alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12 carbon atoms, or alkynyl of 2 to 12 carbon atoms. More typically R^{28} is alkyl of 1 to 8 carbon atoms, alkenyl of 2 to 8 carbon atoms, or alkynyl of 2 to 8 carbon atoms, still more typically, R^{28} is alkyl of 1 to 6 carbon atoms, alkenyl of 2 to 6 carbon atoms, or alkynyl of 2 to 6 carbon atoms. More typically yet, R^{28} is R^{25} .

Each R^{29} is independently R^{22} or R^{28} wherein each R^{28} is substituted with 0 to 3 R^{22} groups. Each of the typical embodiments of R^{28} and R^{22} are typical of R^{29} . More typically R^{29} is substituted with 0, 1, 2, or 3 R^{22} groups.

Each R^{30} is independently H, R^{24} , W or $-R^{24a}W$.


R^4 is $-C(R^{30})_3$, provided that R^4 , taken as a whole, contains 0 to 1 W groups (W is described above) substituted with 0 to 3 R^{29} groups (R^{29} is described above); and, in addition, 1 to 12 carbon atoms substituted with 0 to 3 R^{22} groups (R^{22} is described above). Exemplary embodiments of R^4 are provided as U_1 embodiments in the documents cited in the "Brief Description of Related Art" above.

Typically one R^{30} is H. More typically, one R^{30} is H and the remaining two R^{30} 's are independently R^{24} , W or $-R^{24a}W$. More typically yet, one R^{30} is H, one R^{30} is R^{24} and the remaining R^{30} is independently

R²⁴, W or -R^{24a}W.

In one embodiment of R⁴, one R³⁰ is H, one R³⁰ is R²⁵ and one R³⁰ is R²⁴, W or -R^{24a}W. Typically, one R³⁰ is H and two R³⁰'s are R²⁵. In another embodiment of R⁴, one R³⁰ is H, one R³⁰ is -R^{24a}W and one R³⁰ is R²⁴, W or -R^{24a}W. Typically, one R³⁰ is H, one R³⁰ is -R^{24a}W and one R³⁰ is R²⁴. In another embodiment, one R³⁰ is H and two R³⁰'s are alkyl of 1 to 6 carbon atoms.

In another embodiment, R⁴ is:




wherein R²⁶ is H, -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -OCH₃, -OAc (-O-C(O)CH₃), -OH, -NH₂, or -SH, typically H, -CH₃ or -CH₂CH₃.

Typically each R⁴ (taken as a whole) contains 0-3 W groups each of which is independently substituted with 0-3 R²⁹ groups; and each R⁴ (taken as a whole) in addition contains 1-12 carbon atoms, each carbon atom of which is independently substituted with 0-3 R²² groups. More typically each R⁴ contains 0, 1 or 2 such W groups, more typically yet, 0 or 1 such W group.

In another embodiment, each R³⁰ group (taken as whole) of R⁴ is not so electron withdrawing as to prevent the formation of compound 11. Lowry, T.H. and Richardson, K.S. "Mechanism and Theory in Organic Chemistry" (Harper & Row, 1976) at section 2.2, pages 60-71, and March, J. "Advanced Organic Chemistry" (McGraw-Hill, 1977) at Chapter 9, Quantitative Treatments of the Effect of Structure on Reactivity", pages 251-259, provide details of the electron withdrawing properties of substituent groups. In another embodiment, each R³⁰ group (taken as whole) of R⁴ has a Hammett σ_{para} value of less than about 1, typically less than about 0.75, more typically less than about 0.5. In another embodiment, each R³⁰ group (taken as whole) of R⁴ has a Hammett σ_{para} value of -1.0 to 1.0, more typically -0.75 to 0.75, more typically yet -0.5 to 0.5.

This process embodiment comprises reaction of a compound of the formula:



10

wherein R^{31} is a ketal or acetal, with a lewis acid reagent. Typically R^{31} is -
 $C(R^{30})_2$ - wherein R^{30} is as described above.


Typically, compound 10 is reacted with a Lewis acid catalyst
 5 common in the art, such as $BF_3 \cdot Et_2O$, $TiCl_3$, $TMSOTf$, $SmI_2(THF)_2$,
 $LiClO_4$, $Mg(ClO_4)_2$, $Ln(OTf)_3$ (where $Ln=Yb, Gd, Nd$), $Ti(Oi-Pr)_4$, $AlCl_3$,
 $AlBr_3$, $BeCl_2$, $CdCl_2$, $ZnCl_2$, BF_3 , BCl_3 , BBr_3 , $GaCl_3$, $GaBr_3$, $TiCl_4$, $TiBr_4$,
 $ZrCl_4$, $SnCl_4$, $SnBr_4$, $SbCl_5$, $SbCl_3$, $BiCl_3$, $FeCl_3$, UCl_4 , $ScCl_3$, YCl_3 , $LaCl_3$,
 $CeCl_3$, $PrCl_3$, $NdCl_3$, $SmCl_3$, $EuCl_3$, $GdCl_3$, $TbCl_3$, $LuCl_3$, $DyCl_3$, $HoCl_3$,
 10 $ErCl_3$, $TmCl_3$, $YbCl_3$, ZnI_2 , $Al(OPr^i)_3$, $Al(acac)_3$, $ZnBr_2$, or $SnCl_4$.

Optionally, compound 10 is also treated with a reducing reagent. Typical
 reducing reagents are of the form $B(R^{30})_3$ such as BH_3 . Optionally
 reducing reagents of the form $B(R^{30})_3$ are complexed with common
 solvents such as diethylether and dimethylsulfide. A wide range of borane
 15 reducing reagents are known and will not be described in detail here. For
 example Brown, H.C. "Boranes in Organic Chemistry", (Cornell Univ.
 Press, Ithaca, N.Y., 1972) (Brown) provides a very large number of
 examples such as is found in Part Four, Selective Reductions, pages 209-
 251, Part Five, Hydroboration, pages 255-297, and Part Six, Organoboranes,
 20 pages 301-446.


In a typical embodiment, compound 10 is treated with a lewis acid
 in a nonprotic solvent. More typically, compound 10 is treated with a
 lewis acid and a reducing reagent in a nonprotic solvent.

In a typical embodiment, a solution of 10 in dichlorormethane is
 25 cooled and treated with borane-methyl sulfide complex and trimethylsilyl
 trifluoromethanesulfonate. 10% Aqueous sodium bicarbonate solution is
 slowly added. The mixture is warmed to ambient temperature and
 stirred. The organic layer is filtered and concentrated *in vacuo* to leave
 compound 11. A detailed example of this embodiment is provided as
 30 Example 6 below.

In another example process of this embodiment compound 11 is of the formula:




Another aspect of the present invention is directed to processes for the preparation of compounds of the formula:



wherein:

R^2 , R^4 , R^7 , R^{20} and R^{21} are as defined above.

This process embodiment comprises reaction of a compound of the formula:



with a reducing reagent.


The azide of compound 30 is reduced to form compound 31.

Typically the process comprises treating compound 30 with a reducing agent to form compound 31. More typically the process comprises treating compound 30 with hydrogen gas and a catalyst (such as platinum on carbon or Lindlar's catalyst), or reducing reagents (typically a trisubstituted phosphine such as trialkyl $P(R^{25})_3$ or triaryl phosphine PW_3 , e.g. triphenylphosphine). More typically still, the process comprises treating compound 30 with triphenylphosphine and a base to form


compound 31.

Typically, compound 30 is dissolved in a suitable polar, aprotic solvent such as anhydrous acetonitrile. A solution of anhydrous triphenylphosphine in a suitable solvent such as anhydrous tetrahydrofuran or a mixture of solvents is added dropwise. The mixture is heated at reflux then concentrated *in vacuo* to leave compound 5. A detailed example of this embodiment is provided as Example 9 below.

In another embodiment of this process compound 31 is of the formula:



Another aspect of the present invention is directed to processes for the preparation of compounds of the formula:




wherein:

R^2 , R^4 , R^5 , R^{20} and R^{21} are as described above.

Y^1 is a mono-, di- or unsubstituted amino group. Typically Y^1 is of the formula $-N(R^{30})_2$, a phthalimide or is a nitrogen containing heterocycle (defined above under W), more typically, Y^1 is a phthalimide, more typically yet, a phthalimide salt.

This process embodiment comprises reaction of a compound of the formula:




40

with an amine reagent. Typically, the amine reagent is of the formula HY^1 or a salt of HY^1 , such as, by way of example, NH_3 (McManns, et al., "Bull Soc. Chim. France" 850 (1947)), HY^1 generally (Moussevon, M., et al., "Synth. Commun." 3:177 (1973)) or phthalimide (Gabriel, et al., "Ber." 20:2224 (1887) or Gibson, et al., "Angew. Chem. Int.", 7:919-930 (1968)).


The process comprises treating compound 40 with the amine reagent to produce compound 32. More typically, compound 40 is treated with the amine reagent in a suitable polar aprotic solvent (e.g. CH_3CN , DMF or THF). Optionally compound 40 is treated with the amine reagent and a base. Typical details of this process embodiment can be found in March, "Advanced Organic Chemistry" 4th. ed., pp 425-427.

In another embodiment of this process compound 41 is of the formula:



42

Another aspect of the present invention is directed to processes for the preparation of compounds of the formula:




51

wherein:

R^2 , R^4 , R^{20} , R^{21} and Y^1 are as described above;


This process embodiment comprises reaction of a compound of the formula:



50


with an oxidizing reagent. A wide range of suitable oxidation reagents are common in the art and will not be detailed here. For example House, H.O. "Modern Synthetic Reactions, Second Edition", Chapter 5, pages 259-273, describes the selective oxidation of alcohols. Typical reagents include CrO_3 , $\text{Na}_2\text{Cr}_2\text{O}_7$, KMnO_4 , PDC and PCC. Typical details of this process embodiment can be found in Larock, "Comprehensive Organic Transformations", pp. 604-614; Corey et al., "Tetrahedron Lett." 31:2647-50 (1975); Ley et al., "Chem. Commun" 1625 (1987); Sweon, et al., "J. Org. Chem." 43:2480-2 (1978); and Martin, et al., "J. Org. Chem." 48:4155-56 (1983). Solvents typically include inert polar solvents (e.g. CH_2Cl_2 , toluene or CH_3CN).

In another embodiment of this process compound 51 is of the formula:



52

Another aspect of the present invention is directed to processes for the preparation of compounds of the formula:




61

wherein:

R^2 , R^4 , R^{20} , R^{21} and Y^1 are as described above;

This process embodiment comprises reaction of a compound of the


5 formula:



51


with a base. Typically the base is a hindered amine or hindered alkoxide or the salts of either. More typically the base is of the formula $NaOR^{25}$, KOR^{25} or NR^{25}_3 , more typically yet, DBN, DBU or diisopropyl ethyl amine.

10 In another embodiment of this process compound 61 is of the formula:



62

Another aspect of the present invention is directed to processes for the preparation of compounds of the formula:




71

wherein:

R^2 , R^4 , R^7 , R^{20} , R^{21} and Y^1 are as described above;

This process embodiment comprises reaction of a compound of the

5 formula:



61

with a reductive amination reagent. Typical details of and references to this process embodiment can be found in Larock, *op. cit.*, pp. 421-425.

Another typical description (NaCNBH₃ method) is Borch, "J. Am. Chem.

10 Soc." 93:2897-2904 (1971).

Schemes 1 and 2 depict embodiments of the invention. Detailed descriptions of the processes of Schemes 1 and 2 are provided in the Examples (below).

15 Additional individual process embodiments of the invention include any one or sequential combination of processes AA, AB, AC, AD, AE, AF, AG, AH, AI, AJ, or AK of Schemes 1 and 2. "Sequential combination" as used herein means more than one process wherein the individual processes are performed one after the other in the order shown. Isolation, separation, purification is optionally performed prior to any of


20 the individual processes.


Additional individual process embodiments of the invention include any one or sequential combination of the processes of Example 1,

Example 2, Example 3, Example 4, Example 5, Example 6, Example 7,
Example 8, Example 9, Example 10, Example 11, Example 12 or Example 13.

Scheme 1

5




Scheme 2

- 5 Scheme 3 depicts the synthesis of the neuraminidase inhibitor 206 (R = H₂) by use of alternative nitrogen nucleophiles (March, "Advanced Organic Chemistry" 4th. ed., pp 425-427) to open the epoxide 201. Oxidation of azidoalcohol 202 gives ketone 203 (Larock, "Comprehensive Organic Transformations", pp. 604-614) in which the β-axial NR group
- 10 isomerizes to the α-equatorial configuration 204. Reductive amination of the ketone 204 (Larock, *op. cit.*, pp. 421-425) gives the β-equatorial amine 205 which is acetylated to afford 206. Cleavage of the R moiety (Greene, "Protective Groups in Organic Synthesis", pp. 218-287) gives the
- 15 neuramidase inhibitor 206 (R = H₂).

- 15 Additional individual process embodiments of the invention include any one or sequential combination of processes AL, AM, AN, AO, or AP of Scheme 3.

Scheme 3



Modifications of each of the above schemes leads to various analogs of the specific exemplary materials produced above. The above cited citations describing suitable methods of organic synthesis are applicable to such modifications.

In each of the above exemplary schemes it may be advantageous to separate reaction products from one another and/or from starting materials. The desired products of each step or series of steps is separated and/or purified (hereinafter separated) to the desired degree of homogeneity by the techniques common in the art. Typically such separations involve multiphase extraction, crystallization from a solvent or solvent mixture, distillation, sublimation, or chromatography. Chromatography can involve any number of methods including, for example, size exclusion or ion exchange chromatography, high, medium, or low pressure liquid chromatography, small scale and preparative thin or thick layer chromatography, as well as techniques of small scale thin layer and flash chromatography.

Another class of separation methods involves treatment of a mixture with a reagent selected to bind to or render otherwise separable a desired product, unreacted starting material, reaction by product, or the like. Such reagents include adsorbents or absorbents such as activated

carbon, molecular sieves, ion exchange media, or the like. Alternatively, the reagents can be acids in the case of a basic material, bases in the case of an acidic material, binding reagents such as antibodies, binding proteins, selective chelators such as crown ethers, liquid/liquid ion extraction reagents (LIX), or the like.

Selection of appropriate methods of separation depends on the nature of the materials involved. For example, boiling point, and molecular weight in distillation and sublimation, presence or absence of polar functional groups in chromatography, stability of materials in acidic and basic media in multiphase extraction, and the like. One skilled in the art will apply techniques most likely to achieve the desired separation.

Stereoisomers

The compounds of the invention are enriched or resolved optical isomers at any or all asymmetric atoms. For example, the chiral centers apparent from the depictions are provided as the chiral isomers or racemic mixtures. Both racemic and diastereomeric mixtures, as well as the individual optical isomers isolated or synthesized, substantially free of their enantiomeric or diastereomeric partners, are all within the scope of the invention.

One or more of the following enumerated methods are used to prepare the enantiomerically enriched or pure isomers herein. The methods are listed in approximately their order of preference, i.e., one ordinarily should employ stereospecific synthesis from chiral precursors before chromatographic resolution before spontaneous crystallization.

Stereospecific synthesis is described in the examples. Methods of this type conveniently are used when the appropriate chiral starting material is available and reaction steps are chosen do not result in undesired racemization at chiral sites. One advantage of stereospecific synthesis is that it does not produce undesired enantiomers that must be removed from the final product, thereby lowering overall synthetic yield. In general, those skilled in the art would understand what starting materials and reaction conditions should be used to obtain the desired enantiomerically enriched or pure isomers by stereospecific synthesis. If an unexpected racemization occurs in a method thought to be stereospecific then one needs only to use one of the following separation methods to obtain the desired product.

If a suitable stereospecific synthesis cannot be empirically designed or determined with routine experimentation then those skilled in the art would turn to other methods. One method of general utility is chromatographic resolution of enantiomers on chiral chromatography resins. These resins are packed in columns, commonly called Pirkle columns, and are commercially available. The columns contain a chiral stationary phase. The racemate is placed in solution and loaded onto the column, and thereafter separated by HPLC. See for example, Proceedings Chromatographic Society - International Symposium on Chiral Separations, Sept. 3-4, 1987. Examples of chiral columns that could be used to screen for the optimal separation technique would include Diacel Chriacel OD, Regis Pirkle Covalent Dphenylglycine, Regis Pirkle Type 1A, Astec Cyclobond II, Astec Cyclobond III, Serva Chiral D-DL=Daltosil 100, Bakerbond DNBLau, Sumipax OA-1000, Merck Cellulose Triacetate column, Astec Cyclobond I-Beta, or Regis Pirkle Covalent D-Naphthylalanine. Not all of these columns are likely to be effective with every racemic mixture. However, those skilled in the art understand that a certain amount of routine screening may be required to identify the most effective stationary phase. When using such columns it is desirable to employ embodiments of the compounds of this invention in which the charges are not neutralized, e.g., where acidic functionalities such as carboxyl are not esterified or amidated.

Another method entails converting the enantiomers in the mixture to diastereomers with chiral auxiliaries and then separating the conjugates by ordinary column chromatography. This is a very suitable method, particularly when the embodiment contains free carboxyl, amino or hydroxyl that will form a salt or covalent bond to a chiral auxiliary. Chirally pure amino acids, organic acids or organosulfonic acids are all worthwhile exploring as chiral auxiliaries, all of which are well known in the art. Salts with such auxiliaries can be formed, or they can be covalently (but reversibly) bonded to the functional group. For example, pure D or L amino acids can be used to amidate the carboxyl group of embodiments of this invention and then separated by chromatography.

Enzymatic resolution is another method of potential value. In such methods one prepares covalent derivatives of the enantiomers in the racemic mixture, generally lower alkyl esters (for example of carboxyl), and then exposes the derivative to enzymatic cleavage, generally hydrolysis.

For this method to be successful an enzyme must be chosen that is capable of stereospecific cleavage, so it is frequently necessary to routinely screen several enzymes. If esters are to be cleaved, then one selects a group of esterases, phosphatases, and lipases and determines their activity on the derivative. Typical esterases are from liver, pancreas or other animal organs, and include porcine liver esterase.

If the enantiomeric mixture separates from solution or a melt as a conglomerate, i.e., a mixture of enantiomerically-pure crystals, then the crystals can be mechanically separated, thereby producing the enantiomerically enriched preparation. This method, however, is not practical for large scale preparations and is of no value for true racemic compounds.

Asymmetric synthesis is another technique for achieving enantiomeric enrichment. For example, a chiral protecting group is reacted with the group to be protected and the reaction mixture allowed to equilibrate. If the reaction is enantiomerically specific then the product will be enriched in that enantiomer.

Further guidance in the separation of enantiomeric mixtures can be found, by way of example and not limitation, in "Enantiomers, Racemates, and resolutions", Jean Jacques, Andre Collet, and Samuel H. Wilen (Krieger Publishing Company, Malabar, FL, 1991, ISBN 0-89464-618-4). In particular, Part 2, "Resolution of Enantiomer Mixture", pages 217-435; more particularly, section 4, "Resolution by Direct Crystallization", pages 217-251, section 5, "Formation and Separation of Diastereomers", pages 251-369, section 6, "Crystallization-Induced Asymmetric Transformations", pages 369-378, and section 7, "Experimental Aspects and Art of Resolutions", pages 378-435; still more particularly, section 5.1.4, "Resolution of Alcohols, Transformation of Alcohols into Salt-Forming Derivatives", pages 263-266, section 5.2.3, "Covalent Derivatives of Alcohols, Thiols, and Phenols", pages 332-335, section 5.1.1, "Resolution of Acids", pages 257-259, section 5.1.2, "Resolution of Bases", pages 259-260, section 5.1.3, "Resolution of Amino Acids", page 261-263, section 5.2.1, "Covalent Derivatives of Acids", page 329, section 5.2.2, "Covalent Derivatives of Amines", pages 330-331, section 5.2.4, "Covalent Derivatives of Aldehydes, Ketones, and Sulfoxides", pages 335-339, and section 5.2.7, "Chromatographic Behavior of Covalent Diastereomers", pages 348-354, are cited as examples of the skill of the art.

Salts and Hydrates

The compositions of this invention optionally comprise salts of the compounds herein, for example, Na^+ , Li^+ , K^+ , Ca^{++} and Mg^{++} . Such salts may include those derived by combination of appropriate cations
5 such as alkali and alkaline earth metal ions or ammonium and quaternary amino ions with an acid anion moiety. Monovalent salts are preferred if a water soluble salt is desired.

Metal salts typically are prepared by reacting the metal hydroxide with a compound of this invention. Examples of metal salts which are
10 prepared in this way are salts containing Li^+ , Na^+ , and K^+ . A less soluble metal salt can be precipitated from the solution of a more soluble salt by addition of the suitable metal compound.

In addition, salts may be formed from acid addition of certain organic and inorganic acids, e.g., HCl , HBr , H_2SO_4 , H_3PO_4 or organic
15 sulfonic acids, to basic centers, typically amines. Finally, it is to be understood that the compositions herein comprise compounds of the invention in their un-ionized, as well as zwitterionic form, and combinations with stoichiometric amounts of water as in hydrates.

Also included within the scope of this invention are the salts of the
20 parental compounds with one or more amino acids. Any of the amino acids described above are suitable, especially the naturally-occurring amino acids found as protein components, although the amino acid typically is one bearing a side chain with a basic or acidic group, e.g., lysine, arginine or glutamic acid, or a neutral group such as glycine, serine, threonine,
25 alanine, isoleucine, or leucine.

Additional Uses for the Compounds of This Invention

The compounds of the invention are polyfunctional. As such they represent a unique class of monomers for the synthesis of polymers. By
30 way of example and not limitation, the polymers prepared from the compounds of this invention include polyamides, polyesters and mixed polyester-polyamides.

The present compounds are used as monomers to provide access to polymers having unique pendent functionalities. The compounds of this
35 invention are useful as comonomers with monomers which do not fall within the scope of the invention. Polymers of the compounds of this invention will have utility as cation exchange agents (polyesters or

polyamides) in the preparation of molecular sieves (polyamides), textiles, fibers, films, formed articles and the like. Polymers are prepared by any conventional method, for example, by cross-linking an -OH or -NH₂ group of the compounds of the invention with a diacid comonomer. The preparation of these polymers from the compounds of the invention is conventional per se.

The compounds of the invention are also useful as a unique class of polyfunctional surfactants. Particularly when R⁴ or R² do not contain hydrophilic substituents and are, for example, alkyl, the compounds have the properties of bi-functional surfactants. As such they have useful surfactant, surface coating, emulsion modifying, rheology modifying and surface wetting properties.

As polyfunctional compounds with defined geometry and carrying simultaneously polar and non-polar moieties, the compounds of the invention are useful as a unique class of phase transfer agents. By way of example and not limitation, the compounds of the invention are useful in phase transfer catalysis and liquid/liquid ion extraction (LIX).

The compounds of the invention optionally contain asymmetric carbon atoms. As such, they are a unique class of chiral auxiliaries for use in the synthesis or resolution of other optically active materials. For example, a racemic mixture of carboxylic acids can be resolved into its component enantiomers by: 1) forming a mixture of diastereomeric esters or amides with a compound of the invention containing an -OH or -NH₂ group; 2) separating the diastereomers; and 3) hydrolyzing the ester structure. Further, such a method can be used to resolve the compounds of the invention themselves if optically active acids are used instead of racemic starting materials.

The compounds of this invention are useful as linkers or spacers in preparing affinity absorption matrices, immobilized enzymes for process control, or immunoassay reagents. The compounds herein contain a multiplicity of functional groups that are suitable as sites for cross-linking desired substances. For example, it is conventional to link affinity reagents such as hormones, peptides, antibodies, drugs, and the like to insoluble substrates. These insolubilized reagents are employed in known fashion to absorb binding partners for the affinity reagents from manufactured preparations, diagnostic samples and other impure mixtures. Similarly, immobilized enzymes are used to perform catalytic conversions with facile

recovery of enzyme. Bifunctional compounds are commonly used to link analytes to detectable groups in preparing diagnostic reagents.

Many functional groups in the compounds of this invention are suitable for use in cross-linking. For example, -OH and -NH₂ groups.

5 Suitable protection of reactive groups will be used where necessary while assembling the cross-linked reagent to prevent polymerization of the bifunctional compound of this invention. In general, the compounds here are used by linking them through hydroxyl or amino groups to carboxylic or phosphonic acid groups of the first linked partner, then
10 covalently bonding to the other binding partner through another -OH or -NH₂ group. For example a first binding partner such as a steroid hormone is reacted to form an amide bond with the -NH₂ group of a compound of this invention and then this conjugate is cross-linked through a hydroxyl to cyanogen bromide activated Sepaharose, whereby immobilized steroid
15 is obtained. Other chemistries for conjugation are well known. See for example Maggio, "Enzyme-Immunoassay" (CRC, 1988, pp 71-135) and references cited therein.

The following examples are put forth so as to provide those of
20 ordinary skill in the art with a complete disclosure and description of how to make the compounds and compositions of the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to insure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental
25 errors and deviations should be taken into account. Unless indicated otherwise, parts are parts by weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Examples

30 Example 1

Lactone 100: A solution of quinic acid (20 kg, 104 mol; [α]_D-43.7° (c = 1.12, water); "Merck Index 11th ed", 8071: [α]_D -42° to -44° (water)), 2,2-dimethoxypropane (38.0 kg, 365 mol) and *p*-toluenesulfonic acid monohydrate (200 kg, 1.05 mol) in acetone (80 kg) was heated at reflux for
35 two hours. The reaction was quenched by addition of 21% sodium ethoxide in ethanol (0.340 kg, 1.05 mol) and most of the solvent was distilled *in vacuo*. The residue was partitioned between ethyl acetate (108

kg) and water (30 kg). The aqueous layer was back-extracted with ethyl acetate (13 kg) and the combined organic layers were washed with 5% aqueous sodium bicarbonate (14 kg). Most of the ethyl acetate was distilled *in vacuo* to leave a pale yellow solid residue of 100 which was used directly in the next step.

Example 2

Hydroxy ester 101: A solution of the crude lactone 100 (from 104 mol. (-)-quinic acid) in absolute ethanol (70 kg) was treated with 20% sodium ethoxide in ethanol (0.340kg, 1.05 mol). After two hours at room temperature, acetic acid (0.072 kg, 1.2 mol) was added and the solvent was distilled *in vacuo*. Ethyl acetate (36 kg) was added and the distillation continued to near dryness. The tan solid residue composed of a ca. 5:1 mixture of 101:100 was dissolved in ethyl acetate (9 kg) at reflux and hexane (9 kg) was added. Upon cooling, a white crystalline solid formed which was isolated by filtration to afford a ca. 6.5:1 mixture of 101:100 (19.0 kg, 70% yield).

Example 3

Mesyl ester 102: A solution of a ca. 6.5:1 mixture (18.7 kg, ca. 72 mol) of hydroxy ester 101 and lactone 100 in dichloromethane (77 kg) was cooled to 0-10°C and treated with methanesulfonyl chloride (8.23 kg, 71.8 mol), followed by slow addition of triethylamine (10.1 kg, 100 mol). An additional portion of methanesulfonyl chloride (0.84 kg, 7.3 mol) was added. After one hour, water (10 kg) and 3% hydrochloric acid (11 kg) were added. The layers were separated and the organic layer was washed with water (9 kg), then distilled *in vacuo* to leave a semi-solid residue composed of a ca. 6.5:1 mixture of mesyl ester 102 and mesyl lactone 103. The residue was dissolved in ethyl acetate (11 kg) and cooled to -10° to -20°C for two hours. Mesyl lactone 103 crystallized and was separated by filtration and washed with cold ethyl acetate (11 kg). The filtrate was concentrated to afford mesyl ester 102 as an orange resin (20.5 kg, 84.3% yield).

Example 4

Mesyl acetone 104: A solution of mesyl ester 102 (10.3 kg, 30.4 mol) and pyridine (10.4 kg, 183 mol) in dichloromethane (63 kg) was

cooled to -20° to -30°C and treated portionwise with sulfuryl chloride (6.22 kg, 46 mol). After the exothermic reaction subsided, the resulting slurry was quenched with ethanol (2.4 kg), warmed to 0°C, and washed successively with 16% sulfuric acid (35 kg), water (15 kg) and 5% aqueous sodium bicarbonate (1 kg). The organic layer containing a ca. 4:1:1 mixture of 104:105:106 was concentrated *in vacuo* and ethyl acetate (14 kg) was added. The allylic mesylate 105 was selectively removed by treatment of the ethyl acetate solution with pyrrolidine (2.27 kg, 31.9 mol) and tetrakis(triphenylphosphine)palladium(0) (0.0704 kg, 0.061mol) at ambient temperature for five hours, followed by washing with 16% sulfuric acid (48 kg). The organic layer was filtered through a pad of silica gel (11 kg) and eluted with ethyl acetate (42 kg). The filtrate was concentrated *in vacuo* to leave a thick orange oil composed of a ca. 4:1 mixture of 104:106. The residue was dissolved in ethyl acetate (5.3 kg) at reflux and hexane (5.3 kg) was added. Upon cooling, mesyl acetone 104 crystallized and was separated by filtration and washed with 14% ethyl acetate in hexane (2.1 kg). After drying *in vacuo*, 104 was obtained as pale yellow needles (4.28 kg, 43.4% yield), mp 102-3°C.

20 Example 5

Pentyl ketal 107: A solution of acetone 104 (8.9 kg, 27.8 mol), 3-pentanone (24 kg, 279 mol) and 70% perchloric acid (0.056 kg, 0.39 mol) was stirred for 18 hours. The volatiles were distilled *in vacuo* at ambient temperature and fresh 3-pentanone (30 kg, 348 mol) was added gradually as the distillation progressed. The reaction mixture was filtered, toluene (18 kg) was added, and the resulting solution was washed successively with 6% aqueous sodium bicarbonate (19 kg), water (18 kg) and brine (24 kg). The organic layer was concentrated *in vacuo* and toluene (28 kg) was added gradually as the distillation progressed. When no more distilled, the residual orange oil was composed of pentyl ketal 107 (9.7 kg, 100% yield) and toluene (ca. 2 kg).

Example 6

Pentyl ether 108: A solution of ketal 107 (8.6 kg, 25 mol) in dichloromethane (90 kg) was cooled to -30° to -20°C and treated with borane-methyl sulfide complex (2.1 kg, 27.5 mol) and trimethylsilyl trifluoromethanesulfonate (7.2 kg, 32.5 mol). After one hour, 10%

aqueous sodium bicarbonate solution (40 kg) was slowly added. The mixture was warmed to ambient temperature and stirred for 12 hours. The organic layer was filtered and concentrated *in vacuo* to leave a ca. 8:1 mixture of 108:109 as a gray waxy solid (7.8 kg, 90% yield).

5

Example 7

Epoxide 110: A ca. 8:1 mixture of isomeric pentyl ethers 108:109 (7.8 kg, 22.3 mol) in ethanol (26 kg) was treated with a solution of potassium hydrogen carbonate (3.52 kg, 35 mol) in water (22 kg). After heating at 55°-65°C for two hours, the solution was cooled and twice extracted with hexanes (31 kg, then 22 kg). Unreacted 109 remained in the aqueous ethanol layer. The combined hexane extracts were filtered and concentrated *in vacuo* to leave epoxide 110 as a flocculent white crystalline solid (3.8 kg, 60% yield), mp=54.6°C.

15

Example 8

Hydroxy azide 111: A mixture of epoxide 110 (548 g, 2.0 mol), sodium azide (156 g, 2.4 mol) and ammonium chloride (128.4 g, 2.4 mol) in water (0.265 L) and ethanol (1.065 L) was heated at 70°-75°C for eight hours. Aqueous sodium bicarbonate (0.42 L of 8% solution) was added and the ethanol was distilled *in vacuo*. The aqueous residue was extracted with ethyl acetate (1 L) and the extract was washed with water (0.5 L). The water wash was back-extracted with ethyl acetate (0.5 L). The combined organic extracts were washed with brine (0.5 L), dried over anhydrous sodium sulfate, filtered and concentrated *in vacuo* to leave a ca. 10:1 mixture of isomeric hydroxy azides 111:112 (608 g, 102% yield) as a dark brown oil.

25

Example 9

Aziridine 113: A ca. 10:1 mixture of hydroxy azides 111:112 (608 g, 2.0 mol) was three times co-evaporated *in vacuo* from anhydrous acetonitrile (3 x 0.3 L) and then dissolved in anhydrous acetonitrile (1 L). A solution of anhydrous triphenylphosphine (483 g, 1.84 mol) in anhydrous tetrahydrofuran (0.1 L) and anhydrous acetonitrile (0.92 L) was added dropwise over two hours. The mixture was heated at reflux for six hours then concentrated *in vacuo* to leave a golden paste composed of aziridine 113, triphenylphosphine oxide and traces of triphenylphosphine. The paste was triturated with diethyl ether (0.35 L). Most of the insoluble

35

triphenylphosphine oxide was removed by filtration and washed with diethyl ether (1.5 L). The filtrate was concentrated *in vacuo* to leave a dark brown oil which was dissolved in 20% aqueous methanol and extracted three times with hexanes (3 x 1 L) to remove triphenylphosphine. The hexane extracts were back-extracted with 20% aqueous methanol (0.5 L) and the combined aqueous methanol layers were concentrated *in vacuo*. The residue was twice co-evaporated *in vacuo* from anhydrous acetonitrile (2 x 0.5 L) to leave a dark brown oil composed of aziridine 113 (490 g, 96.8 % yield) and triphenylphosphine oxide (ca. 108 g) which was used directly in the next step.

Example 10

Acetamido azide 115: A mixture of aziridine 113 (490 g, 1.93 mol) and triphenylphosphine oxide (ca. 108 g), sodium azide (151 g, 2.33 mol) and ammonium chloride (125 g, 2.33 mol) in dimethylformamide (1.3 L) was heated at 80°-85°C for five hours. Sodium bicarbonate (32.8 g, 0.39 mol) and water (0.66 L) were added. The amino azide 114 was isolated from the reaction mixture by six extractions with hexanes (6 x 1 L). The combined hexane extracts were concentrated *in vacuo* to ca. 4.5 L total volume and dichloromethane (1.04 L) was added. Aqueous sodium bicarbonate (4.2 L of 8% solution, 3.88 mol) was added, followed by acetic anhydride (198 g, 1.94 mol). After stirring for one hour at ambient temperature, the aqueous layer was discarded. The organic phases were concentrated *in vacuo* to 1.74 kg total weight and dissolved with ethyl acetate (0.209 L) at reflux. Upon cooling, acetamido azide 115 crystallized and was isolated by filtration. After washing with cold 15% ethyl acetate in hexane (1 L) and drying *in vacuo* at ambient temperature, pure 115 was obtained as off-white crystals (361 g, 55% yield), mp 126-132°C.

Example 11

Acetamido amine 116: A mixture of azide 115 (549 g, 1.62 mol) and Lindlar catalyst (50 g) in abs. ethanol (3.25 L) was stirred for eighteen hours while hydrogen (1 atm.) was bubbled through the mixture. Filtration through Celite and concentration of the filtrate *in vacuo* afforded 116 as a foam which solidified on standing (496 g, 98% yield).

Example 12

Phosphate salt of 116: A solution of acetamido amine 116 (5.02 g, 16.1 mmol) in acetone (75 mL) at reflux was treated with 85% phosphoric acid (1.85 g, 16.1 mmol) in abs. ethanol (25 mL). Crystallization commenced immediately and after cooling to 0°C for 12 hours the precipitate was collected by filtration to afford 116•H₃PO₄ as long colorless needles (4.94 g, 75% yield; [α]_D -39.9° (c=1, water)), mp 203-4°C.

Example 13

Hydrochloride salt of 116: A solution of acetamido amine 116 (2.8 g, 8.96 mmol) in abs. ethanol (9 mL) was treated with 2.08 M hydrogen chloride in ethanol (8.6 mL, 17.9 mmol). Most of the ethanol was evaporated *in vacuo* and the oily residue was stirred with ethyl acetate (20 mL) until solid formed. Hexanes (20 mL) were gradually added to the stirred mixture. After one hour at ambient temperature, the solid was collected by filtration, washed with diethyl ether and dried *in vacuo*. This afforded 116•HCl as an off-white solid (2.54 g, 81% yield; [α]_D -43° (c=0.4, water)), mp 206°C.


All literature and patent citations above are hereby expressly incorporated by reference in their entirety at the locations of their citation. Specifically cited sections or pages of the above cited works are incorporated by reference with specificity.

Whenever a compound described herein is substituted with more than one of the same designated group, such as, by way of example and not limitation, "R⁷", "R⁸", "R⁹", "R²⁰", or "R²²", then it will be understood that each of the groups may be the same or different, i.e., each group is independently selected. So for example, the phrase "R²² is" is synonymous with the phrase "each R²² is independently".

The invention has been described in detail sufficient to allow one of ordinary skill in the art to make and use the subject matter of the following claims. It is apparent that certain modifications of the methods and compositions of the following claims can be made within the scope and spirit of the invention.

What is Claimed is:

1. A process for preparation of compounds of the formula:



5

wherein:


R¹ is a cyclic hydroxy protecting group;

R² is a carboxylic acid protecting group;

R³ is a hydroxy protecting group; and

each R²⁰ is independently H or an alkyl of 1 to 12 carbon atoms;

which process comprises reaction of a compound of the formula:




4


with a dehydrating reagent.

2. The process of Claim 1 which further comprises separating compound 5 by treatment with a noble metal complex.

3. The process of Claim 1 wherein compound 5 is of the formula:



4. A process for preparation of compounds of the formula:



5

wherein:

R^2 is a carboxylic acid protecting group;

R^3 is a hydroxy protecting group;

R^4 is $-C(R^{30})_3$;

10

R^5 is H or R^3 ;

R^7 is H or an amino protecting group;

R^8 is H or R^2 ;

R^9 is H or a thiol protecting group;

each R^{20} is independently H or an alkyl of 1 to 12 carbon atoms;

15

each R^{21} is independently R^{20} , Br, Cl, F, I, CN, NO_2 or N_3 ;

each R^{22} is independently F, Cl, Br, I, $-CN$, N_3 , $-NO_2$, $-OR^5$, $-OR^{20}$,

$-N(R^{20})_2$, $-N(R^{20})(R^7)$, $-N(R^7)_2$, $-SR^{20}$, $-SR^9$, $-S(O)R^{20}$, $-S(O)_2R^{20}$, $-S(O)OR^{20}$,

$-S(O)OR^8$, $-S(O)_2OR^{20}$, $-S(O)_2OR^8$, $-C(O)OR^{20}$, $-C(O)OR^8$, $-OC(O)R^{20}$,

$-N(R^{20})(C(O)R^{20})$, $-N(R^7)(C(O)R^{20})$, $-N(R^{20})(C(O)OR^{20})$, $-N(R^7)(C(O)OR^{20})$,

20

$-C(O)N(R^{20})_2$, $-C(O)N(R^7)(R^{20})$, $-C(O)N(R^7)_2$, $-C(NR^{20})(N(R^{20})_2)$,

$-C(N(R^7))(N(R^{20})_2)$, $-C(N(R^{20}))(N(R^{20})(R^7))$, $-C(N(R^7))(N(R^{20})(R^7))$,

$-C(N(R^{20}))(N(R^7)_2)$, $-C(N(R^7))(N(R^7)_2)$, $-N(R^{20})C(N(R^{20}))(N(R^{20})_2)$,

$-N(R^{20})C(N(R^{20}))(N(R^{20})(R^7))$, $-N(R^{20})C(N(R^7))(N(R^{20})_2)$,

$-N(R^7)C(N(R^{20}))(N(R^{20})_2)$, $-N(R^7)C(N(R^7))(N(R^{20})_2)$,

-N(R⁷)C(N(R²⁰))(N(R²⁰)(R⁷)), -N(R²⁰)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R²⁰)C(N(R²⁰))(N(R⁷)₂), -N(R⁷)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R⁷)C(N(R²⁰))(N(R⁷)₂), -N(R²⁰)C(N(R⁷))(N(R⁷)₂),
 -N(R⁷)C(N(R⁷))(N(R⁷)₂), =O, =S, =N(R²⁰), =N(R⁷) or W;

5 R²³ is independently alkyl of 1 to 11 carbon atoms, alkenyl of 2 to 11 carbon atoms, or alkynyl of 2 to 11 carbon atoms;

 R²⁴ is independently R²³ wherein each R²³ is substituted with 0 to 3 R²² groups;

10 R^{24a} is independently alkylene of 1 to 11 carbon atoms, alkenylene of 2 to 11 carbon atoms, or alkynylene of 2-11 carbon atoms any one of which alkylene, alkenylene or alkynylene is substituted with 0-3 R²² groups;


 R²⁸ is independently alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12 carbon atoms, or alkynyl of 2 to 12 carbon atoms;

15 R²⁹ is independently R²² or R²⁸ wherein each R²⁸ is substituted with 0 to 3 R²² groups;

 each R³⁰ is independently H, R²⁴, W or -R^{24a}W; and

 W is carbocycle or heterocycle wherein any one of which carbocycle or heterocycle is substituted with 0 to 3 R²⁹ groups;

20 which process comprises reaction of a compound of the formula:



10

wherein R³¹ is a ketal or acetal, with a lewis acid reagent;


provided that R⁴, taken as a whole, contains:

25 0 to 3 W groups substituted with 0 to 3 R²⁹ groups; and, in addition,

 1 to 12 carbon atoms substituted with 0 to 3 R²² groups.


5. The process of Claim 4 which further comprises treating compound
 30 10 with a reducing reagent.

6. The process of Claim 4 wherein compound 11 is of the formula:



12

7. A process for preparation of compounds of the formula:



31

wherein:

R² is a carboxylic acid protecting group;

R³ is a hydroxy protecting group;

R⁴ is -C(R³⁰)₃;

R⁵ is H or R³;

R⁷ is H or an amino protecting group;

R⁸ is H or R²;

R⁹ is H or a thiol protecting group;

each R²⁰ is independently H or an alkyl of 1 to 12 carbon atoms;

each R²¹ is independently R²⁰, Br, Cl, F, I, CN, NO₂ or N₃;

each R²² is independently F, Cl, Br, I, -CN, N₃, -NO₂, -OR⁵, -OR²⁰,

-N(R²⁰)₂, -N(R²⁰)(R⁷), -N(R⁷)₂, -SR²⁰, -SR⁹, -S(O)R²⁰, -S(O)₂R²⁰, -S(O)OR²⁰,

-S(O)OR⁸, -S(O)₂OR²⁰, -S(O)₂OR⁸, -C(O)OR²⁰, -C(O)OR⁸, -OC(O)R²⁰,

-N(R²⁰)(C(O)R²⁰), -N(R⁷)(C(O)R²⁰), -N(R²⁰)(C(O)OR²⁰), -N(R⁷)(C(O)OR²⁰),

-C(O)N(R²⁰)₂, -C(O)N(R⁷)(R²⁰), -C(O)N(R⁷)₂, -C(NR²⁰)(N(R²⁰)₂),

-C(N(R⁷))(N(R²⁰)₂), -C(N(R²⁰))(N(R²⁰)(R⁷)), -C(N(R⁷))(N(R²⁰)(R⁷)),

-C(N(R²⁰))(N(R⁷)₂), -C(N(R⁷))(N(R⁷)₂), -N(R²⁰)C(N(R²⁰))(N(R²⁰)₂),

-N(R²⁰)C(N(R²⁰))(N(R²⁰)(R⁷)), -N(R²⁰)C(N(R⁷))(N(R²⁰)₂),

-N(R⁷)C(N(R²⁰))(N(R²⁰)₂), -N(R⁷)C(N(R⁷))(N(R²⁰)₂),

-N(R⁷)C(N(R²⁰))(N(R²⁰)(R⁷)), -N(R²⁰)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R²⁰)C(N(R²⁰))(N(R⁷)₂), -N(R⁷)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R⁷)C(N(R²⁰))(N(R⁷)₂), -N(R²⁰)C(N(R⁷))(N(R⁷)₂),
 -N(R⁷)C(N(R⁷))(N(R⁷)₂), =O, =S, =N(R²⁰), =N(R⁷) or W;

5 R²³ is independently alkyl of 1 to 11 carbon atoms, alkenyl of 2 to 11 carbon atoms, or alkynyl of 2 to 11 carbon atoms;

 R²⁴ is independently R²³ wherein each R²³ is substituted with 0 to 3 R²² groups;

10 R^{24a} is independently alkylene of 1 to 11 carbon atoms, alkenylene of 2 to 11 carbon atoms, or alkynylene of 2-11 carbon atoms any one of which alkylene, alkenylene or alkynylene is substituted with 0-3 R²² groups;


 R²⁸ is independently alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12 carbon atoms, or alkynyl of 2 to 12 carbon atoms;

15 R²⁹ is independently R²² or R²⁸ wherein each R²⁸ is substituted with 0 to 3 R²² groups;

 each R³⁰ is independently H, R²⁴, W or -R^{24a}W;

 W is carbocycle or heterocycle wherein any one of which carbocycle or heterocycle is substituted with 0 to 3 R²⁹ groups;

20 which process comprises reaction of a compound of the formula:



30

with a reducing reagent;


provided that R⁴, taken as a whole, contains:

25 0 to 3 W groups substituted with 0 to 3 R²⁹ groups; and, in addition,


1 to 12 carbon atoms substituted with 0 to 3 R²² groups.

8. The process of Claim 7 wherein the reducing reagent is a
 30 trisubstituted phosphine reducing reagent.

9. The process of Claim 7 wherein compound 31 is of the formula:



10. A process for preparation of compounds of the formula:



wherein:

R^2 is a carboxylic acid protecting group;

R^3 is a hydroxy protecting group;

R^4 is $-C(R^{30})_3$;

R^5 is H or R^3 ;

R^7 is H or an amino protecting group;

R^8 is H or R^2 ;

R^9 is H or a thiol protecting group;

each R^{20} is independently H or an alkyl of 1 to 12 carbon atoms;

each R^{21} is independently R^{20} , Br, Cl, F, I, CN, NO_2 or N_3 ;

each R^{22} is independently F, Cl, Br, I, -CN, N_3 , $-NO_2$, $-OR^5$, $-OR^{20}$, $-N(R^{20})_2$, $-N(R^{20})(R^7)$, $-N(R^7)_2$, $-SR^{20}$, $-SR^9$, $-S(O)R^{20}$, $-S(O)_2R^{20}$, $-S(O)OR^{20}$, $-S(O)OR^8$, $-S(O)_2OR^{20}$, $-S(O)_2OR^8$, $-C(O)OR^{20}$, $-C(O)OR^8$, $-OC(O)R^{20}$, $-N(R^{20})(C(O)R^{20})$, $-N(R^7)(C(O)R^{20})$, $-N(R^{20})(C(O)OR^{20})$, $-N(R^7)(C(O)OR^{20})$, $-C(O)N(R^{20})_2$, $-C(O)N(R^7)(R^{20})$, $-C(O)N(R^7)_2$, $-C(NR^{20})(N(R^{20})_2)$, $-C(N(R^7))(N(R^{20})_2)$, $-C(N(R^{20}))(N(R^{20})(R^7))$, $-C(N(R^7))(N(R^{20})(R^7))$, $-C(N(R^{20}))(N(R^7)_2)$, $-C(N(R^7))(N(R^7)_2)$, $-N(R^{20})C(N(R^{20}))(N(R^{20})_2)$, $-N(R^{20})C(N(R^{20}))(N(R^{20})(R^7))$, $-N(R^{20})C(N(R^7))(N(R^{20})_2)$, $-N(R^7)C(N(R^{20}))(N(R^{20})_2)$, $-N(R^7)C(N(R^7))(N(R^{20})_2)$,

-N(R⁷)C(N(R²⁰))(N(R²⁰)(R⁷)), -N(R²⁰)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R²⁰)C(N(R²⁰))(N(R⁷)₂), -N(R⁷)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R⁷)C(N(R²⁰))(N(R⁷)₂), -N(R²⁰)C(N(R⁷))(N(R⁷)₂),
 -N(R⁷)C(N(R⁷))(N(R⁷)₂), =O, =S, =N(R²⁰), =N(R⁷) or W;

5 R²³ is independently alkyl of 1 to 11 carbon atoms, alkenyl of 2 to 11 carbon atoms, or alkynyl of 2 to 11 carbon atoms;

R²⁴ is independently R²³ wherein each R²³ is substituted with 0 to 3 R²² groups;

10 R^{24a} is independently alkylene of 1 to 11 carbon atoms, alkenylene of 2 to 11 carbon atoms, or alkynylene of 2-11 carbon atoms any one of which alkylene, alkenylene or alkynylene is substituted with 0-3 R²² groups;

R²⁸ is independently alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12 carbon atoms, or alkynyl of 2 to 12 carbon atoms;


15 R²⁹ is independently R²² or R²⁸ wherein each R²⁸ is substituted with 0 to 3 R²² groups;

each R³⁰ is independently H, R²⁴, W or -R^{24a}W;

W is carbocycle or heterocycle wherein any one of which carbocycle or heterocycle is substituted with 0 to 3 R²⁹ groups; and

20 Y¹ is a mono-, di- or unsubstituted amino group;

which process comprises reaction of a compound of the formula:



40

with an amine reagent;

provided that R⁴, taken as a whole, contains:


25 0 to 3 W groups substituted with 0 to 3 R²⁹ groups; and, in addition,

1 to 12 carbon atoms substituted with 0 to 3 R²² groups.

11. The process of Claim 10 wherein the amine reagent is a phthalimide reagent.


30

12. The process of Claim 10 wherein compound 41 is of the formula:



42

13. A process for preparation of compounds of the formula:



51

wherein:

R^2 is a carboxylic acid protecting group;

R^3 is a hydroxy protecting group;

R^4 is $-C(R^{30})_3$;

R^5 is H or R^3 ;

R^7 is H or an amino protecting group;

R^8 is H or R^2 ;

R^9 is H or a thiol protecting group;

each R^{20} is independently H or an alkyl of 1 to 12 carbon atoms;

each R^{21} is independently R^{20} , Br, Cl, F, I, CN, NO_2 or N_3 ;

each R^{22} is independently F, Cl, Br, I, -CN, N_3 , $-NO_2$, $-OR^5$, $-OR^{20}$,

$-N(R^{20})_2$, $-N(R^{20})(R^7)$, $-N(R^7)_2$, $-SR^{20}$, $-SR^9$, $-S(O)R^{20}$, $-S(O)_2R^{20}$, $-S(O)OR^{20}$,

$-S(O)OR^8$, $-S(O)_2OR^{20}$, $-S(O)_2OR^8$, $-C(O)OR^{20}$, $-C(O)OR^8$, $-OC(O)R^{20}$,

$-N(R^{20})(C(O)R^{20})$, $-N(R^7)(C(O)R^{20})$, $-N(R^{20})(C(O)OR^{20})$, $-N(R^7)(C(O)OR^{20})$,

$-C(O)N(R^{20})_2$, $-C(O)N(R^7)(R^{20})$, $-C(O)N(R^7)_2$, $-C(NR^{20})(N(R^{20})_2)$,

$-C(N(R^7))(N(R^{20})_2)$, $-C(N(R^{20}))(N(R^{20})(R^7))$, $-C(N(R^7))(N(R^{20})(R^7))$,

$-C(N(R^{20}))(N(R^7)_2)$, $-C(N(R^7))(N(R^7)_2)$, $-N(R^{20})C(N(R^{20}))(N(R^{20})_2)$,

$-N(R^{20})C(N(R^{20}))(N(R^{20})(R^7))$, $-N(R^{20})C(N(R^7))(N(R^{20})_2)$,

$-N(R^7)C(N(R^{20}))(N(R^{20})_2)$, $-N(R^7)C(N(R^7))(N(R^{20})_2)$,

$-N(R^7)C(N(R^{20}))(N(R^{20})(R^7))$, $-N(R^{20})C(N(R^7))(N(R^{20})(R^7))$,

-N(R²⁰)C(N(R²⁰))(N(R⁷)₂), -N(R⁷)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R⁷)C(N(R²⁰))(N(R⁷)₂), -N(R²⁰)C(N(R⁷))(N(R⁷)₂),
 -N(R⁷)C(N(R⁷))(N(R⁷)₂), =O, =S, =N(R²⁰), =N(R⁷) or W;

R²³ is independently alkyl of 1 to 11 carbon atoms, alkenyl of 2 to 11
 5 carbon atoms, or alkynyl of 2 to 11 carbon atoms;

R²⁴ is independently R²³ wherein each R²³ is substituted with 0 to 3
 R²² groups;

R^{24a} is independently alkylene of 1 to 11 carbon atoms, alkenylene
 of 2 to 11 carbon atoms, or alkynylene of 2-11 carbon atoms any one of
 10 which alkylene, alkenylene or alkynylene is substituted with 0-3 R²²
 groups;

R²⁸ is independently alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12
 carbon atoms, or alkynyl of 2 to 12 carbon atoms;


R²⁹ is independently R²² or R²⁸ wherein each R²⁸ is substituted with
 15 0 to 3 R²² groups;

each R³⁰ is independently H, R²⁴, W or -R^{24a}W;

W is carbocycle or heterocycle wherein any one of which carbocycle
 or heterocycle is substituted with 0 to 3 R²⁹ groups; and

Y¹ is a mono-, di- or unsubstituted amino group;

20 which process comprises reaction of a compound of the formula:



50


with an oxidizing reagent;

provided that R⁴, taken as a whole, contains:


25 0 to 3 W groups substituted with 0 to 3 R²⁹ groups; and, in
 addition,

1 to 12 carbon atoms substituted with 0 to 3 R²² groups.

14. The process of Claim 13 wherein compound 51 is of the formula:



5 15. A process for preparation of a compound of the formula:



wherein:

R^2 is a carboxylic acid protecting group;

R^3 is a hydroxy protecting group;

10 R^4 is $-C(R^{30})_3$;

R^5 is H or R^3 ;

R^7 is H or an amino protecting group;

R^8 is H or R^2 ;

R^9 is H or a thiol protecting group;

15 each R^{20} is independently H or an alkyl of 1 to 12 carbon atoms;

each R^{21} is independently R^{20} , Br, Cl, F, I, CN, NO_2 or N_3 ;

each R^{22} is independently F, Cl, Br, I, -CN, N_3 , $-NO_2$, $-OR^5$, $-OR^{20}$,

$-N(R^{20})_2$, $-N(R^{20})(R^7)$, $-N(R^7)_2$, $-SR^{20}$, $-SR^9$, $-S(O)R^{20}$, $-S(O)_2R^{20}$, $-S(O)OR^{20}$,

$-S(O)OR^8$, $-S(O)_2OR^{20}$, $-S(O)_2OR^8$, $-C(O)OR^{20}$, $-C(O)OR^8$, $-OC(O)R^{20}$,

20 $-N(R^{20})(C(O)R^{20})$, $-N(R^7)(C(O)R^{20})$, $-N(R^{20})(C(O)OR^{20})$, $-N(R^7)(C(O)OR^{20})$,

$-C(O)N(R^{20})_2$, $-C(O)N(R^7)(R^{20})$, $-C(O)N(R^7)_2$, $-C(NR^{20})(N(R^{20})_2)$,

$-C(N(R^7))(N(R^{20})_2)$, $-C(N(R^{20}))(N(R^{20})(R^7))$, $-C(N(R^7))(N(R^{20})(R^7))$,

$-C(N(R^{20}))(N(R^7)_2)$, $-C(N(R^7))(N(R^7)_2)$, $-N(R^{20})C(N(R^{20}))(N(R^{20})_2)$,

$-N(R^{20})C(N(R^{20}))(N(R^{20})(R^7))$, $-N(R^{20})C(N(R^7))(N(R^{20})_2)$.

-N(R⁷)C(N(R²⁰))(N(R²⁰)₂), -N(R⁷)C(N(R⁷))(N(R²⁰)₂),
 -N(R⁷)C(N(R²⁰))(N(R²⁰)(R⁷)), -N(R²⁰)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R²⁰)C(N(R²⁰))(N(R⁷)₂), -N(R⁷)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R⁷)C(N(R²⁰))(N(R⁷)₂), -N(R²⁰)C(N(R⁷))(N(R⁷)₂),
 5 -N(R⁷)C(N(R⁷))(N(R⁷)₂), =O, =S, =N(R²⁰), =N(R⁷) or W;

R²³ is independently alkyl of 1 to 11 carbon atoms, alkenyl of 2 to 11 carbon atoms, or alkynyl of 2 to 11 carbon atoms;

R²⁴ is independently R²³ wherein each R²³ is substituted with 0 to 3 R²² groups;

10 R^{24a} is independently alkylene of 1 to 11 carbon atoms, alkenylene of 2 to 11 carbon atoms, or alkynylene of 2-11 carbon atoms any one of which alkylene, alkenylene or alkynylene is substituted with 0-3 R²² groups;

15 R²⁸ is independently alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12 carbon atoms, or alkynyl of 2 to 12 carbon atoms;


R²⁹ is independently R²² or R²⁸ wherein each R²⁸ is substituted with 0 to 3 R²² groups;

each R³⁰ is independently H, R²⁴, W or -R^{24a}W;

20 W is carbocycle or heterocycle wherein any one of which carbocycle or heterocycle is substituted with 0 to 3 R²⁹ groups; and

Y¹ is a mono-, di- or unsubstituted amino group;

which process comprises reaction of a compound of the formula:



51

25 with a base;


provided that R⁴, taken as a whole, contains:

0 to 3 W groups substituted with 0 to 3 R²⁹ groups; and, in addition,


1 to 12 carbon atoms substituted with 0 to 3 R²² groups.

30

16. The process of Claim 15 wherein compound 61 is of the formula:



5 17. A process for preparation of compounds of the formula:



wherein:

R^2 is a carboxylic acid protecting group;

R^3 is a hydroxy protecting group;

10 R^4 is $-C(R^{30})_3$;

R^5 is H or R^3 ;

R^7 is H or an amino protecting group;

R^8 is H or R^2 ;

R^9 is H or a thiol protecting group;

15 each R^{20} is independently H or an alkyl of 1 to 12 carbon atoms;

each R^{21} is independently R^{20} , Br, Cl, F, I, CN, NO_2 or N_3 ;

each R^{22} is independently F, Cl, Br, I, $-CN$, N_3 , $-NO_2$, $-OR^5$, $-OR^{20}$,

$-N(R^{20})_2$, $-N(R^{20})(R^7)$, $-N(R^7)_2$, $-SR^{20}$, $-SR^9$, $-S(O)R^{20}$, $-S(O)_2R^{20}$, $-S(O)OR^{20}$,

$-S(O)OR^8$, $-S(O)_2OR^{20}$, $-S(O)_2OR^8$, $-C(O)OR^{20}$, $-C(O)OR^8$, $-OC(O)R^{20}$,

20 $-N(R^{20})(C(O)R^{20})$, $-N(R^7)(C(O)R^{20})$, $-N(R^{20})(C(O)OR^{20})$, $-N(R^7)(C(O)OR^{20})$,

$-C(O)N(R^{20})_2$, $-C(O)N(R^7)(R^{20})$, $-C(O)N(R^7)_2$, $-C(NR^{20})(N(R^{20})_2)$,

$-C(N(R^7))(N(R^{20})_2)$, $-C(N(R^{20}))(N(R^{20})(R^7))$, $-C(N(R^7))(N(R^{20})(R^7))$,

$-C(N(R^{20}))(N(R^7)_2)$, $-C(N(R^7))(N(R^7)_2)$, $-N(R^{20})C(N(R^{20}))(N(R^{20})_2)$,

$-N(R^{20})C(N(R^{20}))(N(R^{20})(R^7))$, $-N(R^{20})C(N(R^7))(N(R^{20})_2)$,

-N(R⁷)C(N(R²⁰))(N(R²⁰)₂), -N(R⁷)C(N(R⁷))(N(R²⁰)₂),
 -N(R⁷)C(N(R²⁰))(N(R²⁰)(R⁷)), -N(R²⁰)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R²⁰)C(N(R²⁰))(N(R⁷)₂), -N(R⁷)C(N(R⁷))(N(R²⁰)(R⁷)),
 -N(R⁷)C(N(R²⁰))(N(R⁷)₂), -N(R²⁰)C(N(R⁷))(N(R⁷)₂),

5 -N(R⁷)C(N(R⁷))(N(R⁷)₂), =O, =S, =N(R²⁰), =N(R⁷) or W;

R²³ is independently alkyl of 1 to 11 carbon atoms, alkenyl of 2 to 11 carbon atoms, or alkynyl of 2 to 11 carbon atoms;

R²⁴ is independently R²³ wherein each R²³ is substituted with 0 to 3 R²² groups;

10 R^{24a} is independently alkylene of 1 to 11 carbon atoms, alkenylene of 2 to 11 carbon atoms, or alkynylene of 2-11 carbon atoms any one of which alkylene, alkenylene or alkynylene is substituted with 0-3 R²² groups;

15 R²⁸ is independently alkyl of 1 to 12 carbon atoms, alkenyl of 2 to 12 carbon atoms, or alkynyl of 2 to 12 carbon atoms;


R²⁹ is independently R²² or R²⁸ wherein each R²⁸ is substituted with 0 to 3 R²² groups;

each R³⁰ is independently H, R²⁴, W or -R^{24a}W;

20 W is carbocycle or heterocycle wherein any one of which carbocycle or heterocycle is substituted with 0 to 3 R²⁹ groups; and

Y¹ is a mono-, di- or unsubstituted amino group;

which process comprises reaction of a compound of the formula:



61


with a reductive amination reagent;

25 provided that R⁴, taken as a whole, contains:

0 to 3 W groups substituted with 0 to 3 R²⁹ groups; and, in addition,

1 to 12 carbon atoms substituted with 0 to 3 R²² groups.

18. The process of Claim 17 wherein compound 71 is of the formula:



72

19. A process according to any one or sequential combination of processes AA, AB, AC, AD, AE, AF, AG, AH, AI, AJ, or AK of Schemes 1 and 2.
20. A process according to any one or sequential combination of the processes of Example 1, Example 2, Example 3, Example 4, Example 5, Example 6, Example 7, Example 8, Example 9, Example 10, Example 11, Example 12 or Example 13.
21. A process according to any one or sequential combination of processes AL, AM, AN, AO, or AP of Scheme 3.

INTERNATIONAL SEARCH REPORT

Internat. Application No.

PCT/US 97/14813

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07C227/08 C07C247/14 C07D317/46 C07D203/26 C07D303/40
 C07F7/18 C07C227/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C C07D C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	G. ULIBARRI ET AL: "Construction of the bicyclic core structure of the enediyne antibiotic esperamicin-A1 in either enantiomeric form from (-)-quinic acid" JOURNAL OF ORGANIC CHEMISTRY, vol. 60, no. 9, 1995, pages 2753-2761, XP002050553 see page 2755; examples 13,14	1,3
A	L. CHAHOUA ET AL : "Synthesis of (-)-shikimate and (-)-quinic 3-phosphates by differentiation of the hydroxyl functions of (-)-shikimic and (-)-quinic acids" JOURNAL OF ORGANIC CHEMISTRY, vol. 57, no. 21, 1992, pages 5798-5801, XP002050554 see page 5799	4
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 December 1997

Date of mailing of the international search report

14.01.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl
 Fax: (+31-70) 340-3016

Authorized officer

Voyiazoglou, D

INTERNATIONAL SEARCH REPORT

Intern: 1st Application No

PCT/US 97/14813

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	S. FERNANDEZ ET AL: "New and efficient enantiospecific synthesis of (-)-methyl 5-epi-shikimate and methyl 5-epi-quininate from (-)-quinic acid" TETRAHEDRON LETTERS, vol. 38, no. 29, 21 July 1997, pages 5225-5228, XP002050555 see page 5226; examples II,III ---	1
P,X	C. U. KIM ET AL: "influenza neuraminidase inhibitors possessing a novel hydrophobic interaction in the enzyme active site: design, synthesis and structural analysis of carboxylic sialic acid analogues with potent anti-influenza activity" JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol. 119, no. 4, February 1997, pages 681-690, XP002050556 see page 683 - page 685 ---	1-12
P,X	WO 96 26933 A (GILEAD) 6 September 1996 see page 208 - page 214; claim 122 -----	1-12

INTERNATIONAL SEARCH REPORT

Int. l. application No.
PCT/US 97/14813.

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 10-18,21
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
The claims are so broad that for determining the scope of a meaningful international Search due account has been taken of Rule 33.3 PCT; special emphasis was put on the following subject matter: claims 1-9,19-20 ,pages 33-34,41-46

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Information on patent family members

PCT/US 97/14813

Form PCT/ISA/210 (patent family annex) (July 1992)